

AUTOMATIC CONTINUITY OF POLYNOMIAL MAPS AND COCYCLES

TOM MEYEROVITCH AND OMRI NISAN SOLAN

ABSTRACT. By classical theorems of Steinhaus and Weil, any Haar measurable homomorphism between locally compact groups is continuous. In particular, any Lebesgue-measurable homomorphism $\phi : \mathbb{R} \rightarrow \mathbb{R}$ is of the form $\phi(x) = ax$ for some $a \in \mathbb{R}$. In this short note, we prove that any Lebesgue measurable function $\phi : \mathbb{R} \rightarrow \mathbb{R}$ that vanishes under any $d+1$ “difference operators” is a polynomial of degree at most d . More generally, we prove the continuity of any Haar measurable polynomial map between locally compact groups, in the sense of Leibman. We deduce the above result as a direct consequence of a theorem about the automatic continuity of cocycles.

1. STATEMENT OF RESULTS

The Cauchy functional equation

$$\phi(x+y) = \phi(x) + \phi(y) \text{ for all } x, y, \quad (1)$$

is a well-known and fundamental equation in the theory of functional equations. See for instance [8] for a relatively recent survey of some results around the Cauchy functional equation. Around the beginning of the 20th century, it was shown by Steinhaus that any Lebesgue measurable solution $\phi : \mathbb{R} \rightarrow \mathbb{R}$ of the Cauchy functional equation takes the form $\phi(x) = ax$ for some $a \in \mathbb{R}$. Given $t \in \mathbb{R}$ and $f : \mathbb{R} \rightarrow \mathbb{R}$, denote by $\Delta_t f : \mathbb{R} \rightarrow \mathbb{R}$ the function given by

$$\Delta_t f(x) = f(x+t) - f(x).$$

Cauchy’s functional equation can be rewritten as follows:

$$\Delta_x \Delta_y \phi = 0 \text{ for all } x, y.$$

Steinhaus’s theorem can thus be viewed as the case $d = 1$ of the following theorem:

Theorem 1.1. *Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a measurable function. Suppose that there exists $d \geq 0$ that satisfies*

$$\Delta_{t_d} \dots \Delta_{t_0} f(x) = 0 \text{ for all } x, t_0, \dots, t_d \in \mathbb{R}. \quad (2)$$

Then f is a polynomial of degree at most d . Namely, there exists $a_0, \dots, a_d \in \mathbb{R}$ such that

$$f(x) = a_0 + a_1 x + \dots + a_d x^d.$$

By elementary considerations, Steinhaus’s theorem is equivalent to the statement that any Lebesgue measurable homomorphism of \mathbb{R} is continuous. Weil extended this to show that any Haar measurable homomorphism between locally compact Polish groups is continuous. The phenomena of continuity following from apriori weaker assumptions is called “automatic continuity”. See Rosendal’s survey [9] for more on the subject of automatic continuity of homomorphisms.

We will use multiplicative notation for the group operation whenever we do not explicitly assume that the group is commutative, and additive notation whenever we do explicitly

assume that the group is commutative. [Theorem 1.1](#) and Weil's result can be unified using Leibman's notion of *polynomial maps* between groups, which we now recall [\[6\]](#):

Definition 1.2. Let G, H be groups. Given an element $g \in G$, the *difference operator* Δ_g on the space of functions from G to H is given by

$$\Delta_g f(x) = f(gx)f^{-1}(x), \text{ for } f : G \rightarrow H \text{ and } x \in G.$$

In the case of commutative groups G, H , when the group operations are written in additive notation, the difference operation Δ_g with respect to an element $g \in G$ can be written as follows:

$$\Delta_g f(x) = f(x + g) - f(x), \text{ for } f : G \rightarrow H \text{ and } x \in G.$$

Definition 1.3. A *Liebman polynomial of degree at most d* is a function $P : G \rightarrow H$ that vanishes after applying any d difference operators. That is, $P : G \rightarrow H$ is a Liebman map of degree at most d if

$$\Delta_{g_d} \dots \Delta_{g_0} P(x) = 0 \text{ for all } x, g_0, \dots, g_d \in G.$$

Leibman polynomials of degree 0 are precisely constant functions. A Liebman polynomial of degree 1 is an ‘‘affine homomorphism’’, namely function $P : G \rightarrow H$ of the form $P(g) = h\phi(g)$, where $\phi : G \rightarrow H$ is a homomorphism and $h \in H$ is a fixed element.

Theorem 1.4. *Let $P : G \rightarrow H$ be a Haar measurable Liebman polynomial between locally compact Polish groups G, H . Then P is continuous.*

In some cases, there is an explicit description of the (continuous) Liebman polynomials. For instance, it is known that any Liebman polynomial from a group G into a nilpotent group H factors through a nilpotent quotient of G [\[6\]](#). See also [\[3\]](#) for a refinement of this statement and [\[7, Proposition 4.2\]](#) for a quick proof in the case that H is abelian. It is also known that Liebman polynomials between finitely generated nilpotent groups are in fact ordinary polynomials in the Mal'cev coordinates [\[6\]](#). From this, it follows for instance that a real-valued continuous Liebman polynomial from a connected nilpotent group of real matrices is nothing but an ordinary polynomial in the entries of the matrix.

In the case where $G = V$ and $H = W$ are locally compact topological vector spaces over a field of characteristic 0, [Theorem 1.4](#) is a simple corollary of Steinhaus's Theorem.

Proof of Theorem 1.4 when $G = V$ and $H = W$ (sketch) : The proof uses induction on the degree of the polynomial. The degree 0 case is trivial and the degree 1 case is known by Steinhaus's Theorem. Let $P : V \rightarrow W$ be a measurable Liebman polynomial of degree d . We can represent

$$P(x) = \Phi(x, x, x, \dots, x) + P_{d-1}(x)$$

where $P_{d-1} : V \rightarrow W$ is a Liebman polynomial of degree $d - 1$ and $\Phi : V^d \rightarrow W$ is the d -additive Haar measurable map given by $\Phi(x_1, \dots, x_d) = \frac{1}{d!} \Delta_{x_1} \dots \Delta_{x_d} P$. Note that the right-hand side is a constant since P is of degree d . By Steinhaus's Theorem, we deduce that Φ depends continuously on each coordinate, and from here it is not complicated to deduce its continuity. By induction hypothesis, we deduce that P_{d-1} is continuous, and hence P is continuous as well. \square

Using the fact that composition of polynomial maps between nilpotent groups is again polynomial (see [\[6, Prop. 3.22\]](#)) one can use the Mal'cev coordinates parametrization of

Leibman polynomials to deduce the special case of Theorem 1.4 for Leibman polynomials between nilpotent Lie groups, again by applying Steinhaus's Theorem directly.

Obtaining the general case of Theorem 1.4 can seem slightly trickier. However, Theorem 1.4 is an immediate corollary of the following slightly more general result:

Theorem 1.5. *Let $f : G \rightarrow H$ be a Haar measurable function between locally compact Polish groups G, H . If $\Delta_g f : G \rightarrow H$ is continuous for every $g \in G$, then $f : G \rightarrow H$ also continuous.*

Again, Weil's theorem is a particular case because for any homomorphism $\phi : G \rightarrow H$ $\Delta_g \phi : G \rightarrow H$ is a constant function for every $g \in G$, hence continuous. Theorem 1.4 follows directly from Theorem 1.5 by induction on d .

We will introduce one more level of generalization: Let G, H be groups, and let $G \curvearrowright X$ an action of G on a set X . A *cocycle* with respect to this action is a function $c : G \times X \rightarrow H$ that satisfies

$$c(g_1 g_2, x) = c(g_1, g_2 \cdot x) c(g_2, x). \quad (3)$$

Any function $f : X \rightarrow H$ defines a cocycle $\Delta f : G \times X \rightarrow H$ given by

$$\Delta f(g, x) = f(g \cdot x) f^{-1}(x) \text{ for } g \in G, x \in X. \quad (4)$$

The map $f \mapsto \Delta f$ is called the *coboundary map*. It is easy to check that $c = \Delta f$ satisfies (3). A cocycle of the form Δf is called a *coboundary*. For a function $f : G \rightarrow H$, we denote by $\Delta f : G \times G \rightarrow H$ the coboundary associated with the action of G on itself by multiplication from the left.

From Equation (3) it follows that for any cocycle $c : G \times X \rightarrow H$ $c(1_G, x) = 1_H$ for all $x \in X$ and also that for any $(g, x) \in G \times X$ the following holds:

$$c(g^{-1}, x) = (c(g, g^{-1} \cdot x))^{-1}.$$

Theorem 1.6. *Let G, H be locally compact Polish groups, and let X be a locally compact Polish space. Suppose that G acts on X by homeomorphisms, and let $c : G \times X \rightarrow H$ be a Haar measurable cocycle such that $x \mapsto c(g, x)$ is continuous for every $g \in G$. Then $c : G \times X \rightarrow H$ is continuous.*

In that case where $X = \{x_0\}$ is a singleton, any cocycle $c : G \times X \rightarrow H$ is of the form $c(g, x_0) = \phi(g)$, where ϕ is a homomorphism. Thus, the case where X is a singleton, Theorem 1.6 coincides with Weil's theorem on automatic continuity of homomorphisms. Theorem 1.5 follows directly by applying Theorem 1.6 to the coboundary $\Delta f : G \times G \rightarrow H$.

Somewhat different aspects of continuity for cocycles on topological groups have been explored in the literature. In [1] Becker has shown that whenever G, H are Polish groups and X is a Polish space, any Borel measurable cocycle $c : G \times X \rightarrow H$ is *nearly essentially continuous* in the sense that for any Borel probability measure on X which is G -quasi-invariant and any $\epsilon > 0$ there exists a compact set $K \subseteq X$ such that $\mu(K) > 1 - \epsilon$ and a Borel subset $S \subseteq G \times X$ such that $(g, x) \in S$ whenever $x, gx \in K$ and so that the restriction of c to K is continuous [1, Theorem 1.4.5]. Several other results regarding the continuity of cocycles are discussed in [1] and earlier references within. As far as we are aware, Theorem 1.6 is not directly covered by existing results in the literature.

We remark that in this paper we do not explore possible extensions beyond the setup of locally compact Polish groups. Automatic continuity holds in the context of Haar measurable

homomorphisms between (not necessarily Polish) locally compact groups [5, 4]. Answering a longstanding problem originating in Christensen's seminal work on Haar null sets, Rosendal proved that any universally measurable homomorphism between Polish groups is automatically continuous [10]. Rosendal's proof utilizes a "Steinhaus type principle" due to Christensen [2]. One can wonder if Theorem 1.6 can be similarly extended beyond the setting of locally compact Polish groups.

Acknowledgements: The authors thank Nachi Avraham Re'em, Uri Bader, Eli Glasner, Yair Glasner, and the anonymous referee for useful references, historical remarks and helpful comments. The first author is supported by the Israel Science Foundation grant no. 985/23. The second author is supported by ERC 2020 grant HomDyn (grant no. 833423).

2. CONTINUITY OF COCYCLES

We proceed to prove Theorem 1.6. The proof is based on the following theorem of André Weil, which is a generalization of Steinhaus's theorem from the case $G = \mathbb{R}$ (see for instance [9, Theorem 2.3]):

Theorem 2.1 (H. Stienhaus- A. Weil). *Let G be a locally compact Polish group and $E \subseteq G$ a Haar measurable set of positive Haar measure. Then $EE^{-1} \subseteq G$ contains an open neighborhood of 1_G .*

We first introduce a simple measure-theoretic lemma that does not involve groups:

Lemma 2.2. *Let Z, Y be separable metric spaces, with Z locally compact. Let μ be a Radon measure on Z and let $F : Z \times Z \rightarrow Y$ be a measurable function such that $x_2 \mapsto F(x_1, x_2)$ is continuous for every $x_1 \in Z$. Then for any $\epsilon > 0$ and any measurable set $E_0 \subseteq Z$ with $\mu(E_0) > 0$, there exists a measurable set $E \subseteq E_0$ with $\mu(E) > 0$ such that the diameter of the image of $E \times E$ under F is at most ϵ in the following sense:*

$$d_Y(F(x_1, x_2), F(\tilde{x}_1, \tilde{x}_2)) < \epsilon \text{ for every } x_1, x_2, \tilde{x}_1, \tilde{x}_2 \in E.$$

Proof. Let Z, μ and $F : Z \times Z \rightarrow Y$ be as in the statement. Fix $\epsilon > 0$ and a measurable set $E_0 \subseteq Z$ with $\mu(E_0) > 0$. Because μ is a Radon measure, we can find a compact set $K \subseteq E_0$ such that $\mu(K) > 0$. Because K is a compact metric space, it follows that the space $C(K, Y)$ of continuous functions from K to Y is separable. Given $f \in C(K, Y)$ and $\delta > 0$, let

$$A_{f,\delta} = \left\{ \tilde{f} \in C(K, Y) : d_Y(\tilde{f}(x), f(x)) < \delta \forall x \in K \right\},$$

and let

$$E_{f,\delta} = \{x \in K : F_x \in A_{f,\delta}\}.$$

where for $x \in Z$ $F_x \in C(Z, Y)$ is given by $F_x(\tilde{x}) = F(x, \tilde{x})$. Choose $\delta = \epsilon/4$.

Then for every $f \in C(K, Y)$ and $\delta > 0$ the set $A_{f,\delta} \subseteq C(K, Y)$ is an open neighborhood of f . Since $C(K, Y)$ is separable there is a countable collection $T \subseteq C(K, Y)$ such that $C(K, Y) = \bigcup_{f \in T} A_{f,\delta}$. Hence $K = \bigcup_{f \in T} E_{f,\delta}$. Because $\mu(K) > 0$, there exists $f \in T$ such that $\mu(E_{f,\delta}) > 0$. Fix such f . Since f is continuous and K is compact, there exists a finite open cover \mathcal{U} of K such that for every $U \in \mathcal{U}$ and every $x_1, x_2 \in U$ we have $d_Y(f(x_1), f(x_2)) < \delta$. Because $\mu(E_{f,\delta}) > 0$ and \mathcal{U} covers K , there exists $U \in \mathcal{U}$ such that $\mu(E_{f,\delta} \cap U) > 0$. Choose $E = E_{f,\delta} \cap U$. Clearly, $E \subseteq E_0$ is measurable and $\mu(E) > 0$. Now for any $x_1, x_2, \tilde{x}_1, \tilde{x}_2 \in E$, we have

$$d_Y(F(x_1, x_2), F(\tilde{x}_1, \tilde{x}_2)) \leq d_Y(F(x_1, x_2), f(x_2)) + d_Y(f(x_2), f(\tilde{x}_2)) + d_Y(f(\tilde{x}_2), F(\tilde{x}_1, \tilde{x}_2)).$$

Since $x_1, x_2 \in E \subseteq E_{f,\delta}$ and $x_2, \tilde{x}_2 \in K$ we have that $d_Y(F(x_1, x_2), f(x_2)) < \delta$ and $d_Y(f(\tilde{x}_2), F(\tilde{x}_1, \tilde{x}_2)) < \delta$. Since $x_2, \tilde{x}_2 \in E \subseteq U$, we have that $d_Y(f(x_2), f(\tilde{x}_2)) < \delta$. So altogether, $d_Y(F(x_1, x_2), F(\tilde{x}_1, \tilde{x}_2)) < \delta$. \square

Proof of Theorem 1.6. Let $c : G \times X \rightarrow H$ be a Haar measurable cocycle such that $(g, x) \mapsto c(g, x)$ is continuous with respect to the variable $x \in X$ for every $g \in G$. We first prove that for every $x_0 \in X$ the function $c : G \times X \rightarrow H$ is continuous at $(1_G, x_0)$. Equivalently, we need to prove the following: For every $x_0 \in X$ and every open neighborhood $V \subseteq H$ of 1_H there exist open sets $U \subseteq G$ and $W \subseteq X$ with $(1_G, x_0) \in U \times W$ so that $U \times W \subseteq c^{-1}(V)$.

Fix $x_0 \in X$ and any open set $V \subseteq H$ with $1_H \in V$. Find $\epsilon > 0$ such that V contains the ball of radius ϵ around 1_H .

Since X is locally compact, we can find a compact subset $X_0 \subseteq X$ that contains an open neighborhood of x_0 . For every $g \in G$ there exists $r_g > 0$ so that $c(g, x)$ is contained in the ball of radius r_g in H around 1_H for every $x \in X_0$. Thus there exists a compact subset $K_1 \subseteq H$ such that the set

$$E_0 = \{g \in G : c(g, x) \in K_1 \text{ for all } x \in X_0\}.$$

has a positive Haar measure.

Since $K_1 \subseteq H$ is compact, there exists $\delta > 0$ such that for any ball $B \subseteq K_1$ of radius δ , the set BB^{-1} is contained in a ball of radius ϵ around 1_H in H . Set $Y = H$ and $Z = G$. Define $F : Z \times Z \rightarrow Y$ by $F(g_1, g_2) = c(g_1, g_2(x_0))$. Then we can apply Lemma 2.2 and find a set $E_1 \subseteq E_0$ of positive Haar measure and a ball \tilde{B} of radius less than δ in K_1 such that $c(g_1, g_2(x_0)) \in \tilde{B}$ for every $g_1, g_2 \in E_1$. Since c is continuous in the second argument, we can choose a slightly larger ball B with $\tilde{B} \subseteq B$ such that B still has radius at most δ and for every $g_1 \in E_1$ we can find an open neighborhood W_{g_1} of x_0 such that $c(g_1, g_2(x)) \in \tilde{B}$ for every $x \in W_{g_1}$ and every $g_2 \in E_1$. We can thus find a measurable subset $E \subseteq E_1$ of positive Haar measure and an open neighborhood W of x_0 such that for every $x \in W$ and every $g_1, g_2 \in E$ we have $c(g_1, g_2(x)) \in B$.

Let $U = EE^{-1} \subseteq G$. By Weil's Theorem 2.1, U contains an open neighborhood of 1_G . Now suppose $g \in U$ and $x \in W$. Then there exists $g_1, g_2 \in E$ such that $g = g_1 g_2^{-1}$, and so for every $x \in W$ we have

$$c(g, x) = c(g_1 g_2^{-1}, x) = c(g_1, g_2^{-1} \cdot x) (c(g_2, g_2^{-1} \cdot x))^{-1} \in BB^{-1} \subseteq V.$$

This completes the proof that $c : G \times X \rightarrow H$ is continuous at $(1_G, x_0)$ for every $x_0 \in X$. To complete the proof that $c : G \times X \rightarrow H$ is continuous at every point $(g_0, x_0) \in G \times X$ recall the cocycle equation

$$c(gg_0, x) = c(g, g_0 \cdot x) c(g_0, x).$$

So

$$\begin{aligned} \lim_{(g,x) \rightarrow (g_0, x_0)} c(g, x) &= \lim_{(g,x) \rightarrow (1_G, x_0)} c(gg_0, x) = \\ \lim_{(g,x) \rightarrow (1_G, x_0)} c(g, g_0 \cdot x) c(g_0, x) &= \lim_{(g,x) \rightarrow (1_G, x_0)} c(g, g_0 \cdot x) \cdot \lim_{(g,x) \rightarrow (1_G, x_0)} c(g_0, x) = 1_H \cdot c(g_0, x_0). \end{aligned}$$

\square

REFERENCES

1. Howard Becker, *Cocycles and continuity*, Trans. Amer. Math. Soc. **365** (2013), no. 2, 671–719. MR 2995370
2. Jens Peter Reus Christensen, *Borel structures in groups and semigroups*, Math. Scand. **28** (1971), 124–128. MR 308322
3. Ben Green and Terence Tao, *The quantitative behaviour of polynomial orbits on nilmanifolds*, Ann. of Math. (2) **175** (2012), no. 2, 465–540. MR 2877065
4. Adam Kleppner, *Correction to: “Measurable homomorphisms of locally compact groups”* [Proc. Amer. Math. Soc. **106** (1989), no. 2, 391–395; MR0948154 (89k:22005)], Proc. Amer. Math. Soc. **111** (1991), no. 4, 1199–1200. MR 1069290
5. Yulia Kuznetsova, *On continuity of measurable group representations and homomorphisms*, Studia Math. **210** (2012), no. 3, 197–208. MR 2983459
6. A. Leibman, *Polynomial mappings of groups*, Israel J. Math. **129** (2002), 29–60. MR 1910931
7. Tom Meyerovitch, Idan Perl, Matthew Tointon, and Ariel Yadin, *Polynomials and harmonic functions on discrete groups*, Trans. Amer. Math. Soc. **369** (2017), no. 3, 2205–2229. MR 3581232
8. Daniel Reem, *Remarks on the Cauchy functional equation and variations of it*, Aequationes Math. **91** (2017), no. 2, 237–264. MR 3627381
9. Christian Rosendal, *Automatic continuity of group homomorphisms*, Bull. Symbolic Logic **15** (2009), no. 2, 184–214. MR 2535429
10. ———, *Continuity of universally measurable homomorphisms*, Forum Math. Pi **7** (2019), e5, 20. MR 3996719

BEN GURION UNIVERSITY OF THE NEGEV. DEPARTEMENT OF MATHEMATICS. BE’ER SHEVA, 8410501, ISRAEL

THE HEBREW UNIVERSITY OF JERUSALEM. EINSTEIN INSTITUTE OF MATHEMATICS EDMOND J. SAFRA CAMPUS GIVAT RAM. JERUSALEM, 9190401, ISRAEL