Some exact constants for bilateral approximations in quasi-normed groups

Oleh Lopushansky

Abstract

We establish inverse and direct theorems on best approximations in quasi-normed Abelian groups in the form of bilateral Bernstein-Jackson inequalities with exact constants. Using integral representations for quasi-norms of functions \boldsymbol{f} in Lebesgue's spaces by decreasing rearrangements $\boldsymbol{f^*}$ with the help of approximation \boldsymbol{E} -functionals, error estimates are found. Examples of numerical calculations, spectral approximations of self-adjoint operators obtained by obtained estimates are given.

Keywords: Bernstein-Jackson inequalities, quasi-normed Abelian groups, exact approximation constants

MSC Classification: 42B35, 41A44, 41A17, 46B70

1 Introduction and main results

This article is devoted to the proof of a bilateral version of Bernstein-Jackson inequalities for quasi-normed Abelian groups. The case of similar inequalities on Gaussian Hilbert spaces of random variables was analyzed in the previous publication [10]. In the studied case, we use the approximation scale

$$\mathcal{B}_{\tau}^{s}(\mathfrak{A}_{0},\mathfrak{A}_{1})=(\mathfrak{A}_{0},\mathfrak{A}_{1})_{\vartheta,q}^{1/\vartheta}\,,\quad s+1=1/\vartheta,\quad \tau=\vartheta q$$

of compatible couple of quasi-normed Abelian groups with $0 < \vartheta < 1$ and $0 < q \le \infty$, defined by the Lions-Peetre method of real interpolation. It should be noted that in partial cases this scale coincides with the known scale of Besov spaces (see [1, Thm 7.2.4], [5, 6],[14, Thm 2.5.4]).

We also analyze the problem of accurate error estimates in quasi-normed spaces $L^p_\mu = L^p_\mu(\mathfrak{A})$ ($0 \le p \le \infty$) of \mathfrak{A} -valued functions $f \colon X \to \mathfrak{A}$ on a measure

⁰mailto:olopuszanski@ur.edu.pl Institute of Mathematics, University of Rzeszów, Poland

2

space (X, μ) . Using the decreasing rearrangement method (see e.g. [3, 4]), we reduce this problem to the simpler case for uniquely defined decreasing functions f^* on $(0, \infty)$.

The scale of quasi-normed spaces $\mathcal{B}^s_{\tau}(\mathfrak{A}_0,\mathfrak{A}_1)$ endowed with the quasinorm $\|\cdot\|_{\mathcal{B}^s_{\tau}}$ is determined by the best approximation E-functional

$$E(t, a; \mathfrak{A}_0, \mathfrak{A}_1) = \inf\{|a - a_0|_1 : |a_0|_0 < t\}, \quad a \in \mathfrak{A}_1$$

which fully characterizes the accuracy of error estimates (see e.g. [1, 11, 15]). In Theorem 1 we establish the bilateral version of Bernstein-Jackson inequalities for quasi-normed groups with the exact constants $c_{s,\tau}$ depend only on basis parameters $\mathcal{B}_{\tau}^{s}(\mathfrak{A}_{0}, \mathfrak{A}_{1})$

$$t^s E(t, a) \le c_{s,\tau} |a|_{\mathcal{B}^s_{\tau}} \le 2^{1/2} |a|_0^s |a|_1, \quad c_{s,\tau} = \left[\frac{s}{\tau(s+1)^2}\right]^{1/\tau}$$

for all $a \in \mathfrak{A}_0 \cap \mathfrak{A}_1$ and $t, \tau > 0$. The left inequality allows the unique extension to the whole approximation spaces $\mathcal{B}^s_{\tau}(\mathfrak{A}_0, \mathfrak{A}_1)$ in form of the Jackson-type inequality

$$E(t, a) \le c_{s,\tau} t^{-s} |a|_{\mathcal{B}^s_{\tau}}$$
 for all $a \in \mathcal{B}^s_{\tau}(\mathfrak{A}_0, \mathfrak{A}_1)$.

In Section 3 these results are applied to the quasi-normed Abelian groups $L^p_\mu = L^p_\mu(\mathfrak{A})$ with $0 \le p \le \infty$ and a positive Radon measure μ on a measure space X of measurable functions $f: X \to \mathfrak{A}$ endowed with the κ -norm

$$||f||_p = \begin{cases} \left(\int |f(x)|^p \, \mu(dx) \right)^{1/p} & \text{if } 0$$

where supp $f \subset X$ is a measurable subset such that $f|_{X\setminus \text{supp }f} = 0$ and $f \neq 0$ almost everywhere on supp f with respect to the measure μ . In Theorem 7 the equality

$$f^*(t) = E\big(t,f;\; L_\mu^0,L_\mu^\infty\big), \quad f \in L_\mu^\infty,$$

where f^* is the decreasing rearrangement of \mathfrak{A} -valued functions $f \in L^p_\mu$ and, as a consequence, the following equalities

$$||f||_{\tau/s} = \begin{cases} \left(\int_0^\infty \left[t^s f^*(t) \right]^\tau \frac{dt}{t} \right)^{1/\tau} & \text{if } \tau < \infty \\ \sup_{0 < t < \infty} t^s f^*(t) & \text{if } \tau = \infty, \end{cases}$$

are established. We also compute exact constants in the Jackson-type inequalities

$$f^*(t) < t^s 2^{(s+1)/2} c_{s,\tau} ||f||_{\tau/s}$$
 for all $f \in L^{\tau/s}$

that estimate measurable errors for decreasing rearrangements f^* , as well as the bilateral Bernstein-Jackson inequalities

$$t^{-s}2^{-(s+1)/2}f^*(t) \le c_{s,\tau} \|f\|_{\tau/s} \le \|f\|_0^s \|f\|_{\infty}$$
 for all $f \in L^0 \cap L^{\infty}$, where

$$2^{-1/2\theta}C_{\theta,q} = \begin{cases} c_{s,\tau} & \text{if } \tau < \infty \\ 1 & \text{if } \tau = \infty. \end{cases}$$

In particular, the following inequality holds,

$$f^*(t) \le \frac{2}{\pi t} \int |f| \mu(dx)$$
 for all $f \in L^1_\mu$.

The numerical algorithm resulting from the Theorem 7 was carried out on the example of inverse Gaussian distribution. Examples of applications for accurate estimates of spectral approximations of self-adjoint operators are given in Section 4.

Note that basic notations used in this work can be found in [1, 18].

2 Best approximation scales of quasi-normed Abelian groups

In what follows, we study the κ -normed Abelian groups $(\mathfrak{A}, |\cdot|)$ relative to an operation "+", where κ -norm $|\cdot|$ with the constant $\kappa \geq 1$ is determined by the assumptions (see e.g. [1, 11]):

```
|0| = 0 and |a| > 0 for all nonzero a \in \mathfrak{A},

|a| = |-a| for all a \in \mathfrak{A},

|a+b| \le \kappa(|a|+|b|) with \kappa independent on a,b \in \mathfrak{A}
```

As is known, each κ -norm $|\cdot|$ can be replaced by an equivalent 1-norm $|\cdot|'$ such that $|\cdot|' \leq |\cdot|^{\rho} \leq 2|\cdot|'$ by taking $(2\kappa)^{\rho} = 2$ with a suitable $\rho > 0$ (see e.g. [15]). We will not assume the completeness of groups.

Given a compatible couple $(\mathfrak{A}_0, \mathfrak{A}_1)$ of κ_i -normed groups $(\mathfrak{A}_i, |\cdot|_i)$ with $i = \{0, 1\}$ and $a = a_0 + a_1$ in the sum $\mathfrak{A}_0 + \mathfrak{A}_1$ such that $a_i \in \mathfrak{A}_i$ we define the best approximation E-functional $E(t, a; \mathfrak{A}_0, \mathfrak{A}_1)$ with $a \in \mathfrak{A}_0 + \mathfrak{A}_1$ and t > 0 in the form (see [1, no.7])

$$E(t, a) = E(t, a; \mathfrak{A}_0, \mathfrak{A}_1) = \inf\{|a - a_0|_1 : |a_0|_0 < t\}, \quad a \in \mathfrak{A}_1.$$
 (1)

4

Definition 1 For any $\tau = q\vartheta$ and $s+1 = 1/\vartheta$ with $0 < \vartheta < 1$ and $0 < q \le \infty$ the κ -normed best approximation scales of Abelian groups are defined to be

$$\mathcal{B}_{\tau}^{s}(\mathfrak{A}_{0},\mathfrak{A}_{1}) = \left\{ a \in \mathfrak{A}_{0} + \mathfrak{A}_{1} \colon |a|_{\mathcal{B}_{\tau}^{s}} < \infty \right\},$$

$$|a|_{\mathcal{B}_{\tau}^{s}} = \begin{cases} \left(\int_{0}^{\infty} \left[t^{s} E(t, a) \right]^{\tau} \frac{dt}{t} \right)^{1/\tau} & \text{if } \tau < \infty \\ \sup_{0 < t < \infty} t^{s} E(t, a) & \text{if } \tau = \infty, \end{cases}$$

$$(2)$$

where (by [1, Lemma 7.1.6]) $\kappa = 2\kappa_1 \max(\kappa_0^s, \kappa_1^s) \max(1, 2^{-1/q'}) \max(1, 2^{s-1}), (1/q' = 1 - 1/q).$

Given a compatible couple $(\mathfrak{A}_0, \mathfrak{A}_1)$, we also define the quadratic functional of Lions-Peetre's type (see e.g. [12, 15])

$$K_2(t,a) = K_2(t,a; \mathfrak{A}_0, \mathfrak{A}_1) = \inf_{a=a_0+a_1} (|a_0|_0^2 + t^2|a_1|_1^2)^{1/2}, \quad t > 0.$$

We will use the quadratically modified real interpolation method. Define the interpolation Abelian group of Lions-Peetre's type $(\mathfrak{A}_0,\mathfrak{A}_1)_{\vartheta,q}$ endowed with the quasinorm $\|\cdot\|_{(\mathfrak{A}_0,\mathfrak{A}_1)_{\vartheta,q}}$,

$$K_{\theta,q}\left(\mathfrak{A}_{0},\mathfrak{A}_{1}\right)=\left(\mathfrak{A}_{0},\mathfrak{A}_{1}\right)_{\vartheta,q}=\left\{a\in\mathfrak{A}_{0}+\mathfrak{A}_{1}\right\},$$

$$|a|_{\left(\mathfrak{A}_{0},\mathfrak{A}_{1}\right)_{\vartheta,q}}=\left\{\left(\int_{0}^{\infty}\left[t^{-\vartheta}K_{2}(t,a)\right]^{q}\frac{dt}{t}\right)^{1/q}\text{ if }q<\infty\right.$$

$$\sup_{0< t<\infty}t^{-\vartheta}K_{2}(t,a)\qquad\text{if }q=\infty.$$

Following [7, 10], we extend the use of approximation constants $c_{s,\tau}$ to the case of arbitrary quasi-normed Abelian groups $\mathcal{B}_{\tau}^{s}(\mathfrak{A}_{0},\mathfrak{A}_{1})$ described in the classic works [15, 16], where

$$c_{s,\tau} := \left\{ \begin{bmatrix} \frac{s}{\tau(s+1)^2} \end{bmatrix}^{1/\tau} & \text{if } \tau < \infty \\ 1 & \text{if } \tau = \infty \end{cases}$$
 (3)

is determined by the normalization factor $N_{\vartheta,q}$ from [1, Thm 3.4.1] to be

$$c_{s,\tau} = N_{\vartheta,q}^{1/q} \left(\vartheta q^2\right)^{-1/q\vartheta}, \quad N_{\vartheta,q} := \left[q\vartheta(1-\vartheta)\right]^{1/q}. \tag{4}$$

The following theorem establishes the bilateral form of approximation inequalities with exact constants for the case of quasi-normed Abelian groups.

Theorem 1 The bilateral Bernstein-Jackson inequalities with the exact constant $c_{s,\tau}$

$$t^{s}E(t,a) \le c_{s,\tau}|a|_{\mathcal{B}^{s}_{-}} \le 2^{1/2}|a|_{0}^{s}|a|_{1} \quad for \ all \quad a \in \mathfrak{A}_{0} \cap \mathfrak{A}_{1}$$
 (5)

hold. There is a unique extension of Jackson's inequality on the whole approximation scale

$$E(t,a) \le t^{-s} c_{s,\tau} |a|_{\mathcal{B}^s_{\tau}} \quad \text{for all} \quad a \in \mathcal{B}^s_{\tau}(\mathfrak{A}_0,\mathfrak{A}_1), \quad r > 0.$$
 (6)

Proof Let $0 < q < \infty$ and $\alpha = |a|_{\mathfrak{A}_1}/|a|_{\mathfrak{A}_0}$ with a nonzero $a \in \mathfrak{A}_0 \cap \mathfrak{A}_1$. Since

$$K_2(t,a)^2 = \inf_{a=a_0+a_1} \left(|a_0|_0^2 + t^2 |a_1|_1^2 \right) \le |a|_0^2 \min(1,\alpha^2 t^2) = \min\left(|a|_0^2, t^2 |a|_1^2 \right)$$

or otherwise $K_2(t,a) \leq |a|_0 \min(1,\alpha t) = \min(|a|_0,t|a|_1)$, we get the inequality

$$|a|_{(\mathfrak{A}_0,\mathfrak{A}_1)_{\vartheta,q}}^q \le |a|_1^q \int_0^\alpha t^{-1+q(1-\vartheta)} dt + |a|_0^q \int_\alpha^\infty t^{-1-\vartheta q} dt.$$

Calculating the integrals and using that $\alpha = |a|_1/|a|_0$, we obtain

$$|a|_{(\mathfrak{A}_0,\mathfrak{A}_1)_{\vartheta,q}}^q \leq \frac{\alpha^{q(1-\vartheta)}}{q(1-\vartheta)}|a|_1^q + \frac{\alpha^{-\vartheta q}}{\vartheta q}|a|_0^q = \frac{1}{q\vartheta(1-\vartheta)}\left(|a|_0^{1-\vartheta}|a|_1^\vartheta\right)^q.$$

Let it be now $q = \infty$. Then the inequality

$$t^{-\vartheta}K_2(t,a) \le \min\left(t^{-\vartheta}|a|_0, t^{1-\vartheta}|a|_1\right)$$

holds. Taking in this case $t = |a|_0/|a|_1$, we obtain that $t^{-\vartheta}K_2(t,a) \leq |a|_0^{1-\vartheta}|a|_1^{\vartheta}$. Combining previous inequalities and taking into account (4), we have

$$||a||_{(\mathfrak{A}_0,\mathfrak{A}_1)_{\vartheta,q}} \le \begin{cases} N_{\vartheta,q}^{-1}|a|_0^{1-\vartheta}|a|_1^{\vartheta} & \text{if } q < \infty \\ |a|_0^{1-\vartheta}|a|_1^{\vartheta} & \text{if } q = \infty \end{cases}, \quad a \in \mathfrak{A}_0 \cap \mathfrak{A}_1. \tag{7}$$

For further considerations, we need the functional

$$K_{\infty}(v,a) := \inf_{a=a_0+a_1} \max \left(|a_0|_{\mathfrak{A}_0}, v|a_1|_{\mathfrak{A}_1} \right).$$

Now we will use the known properties of the considered functionals that

$$v^{-\theta}K_{\infty}(v,f) \to 0$$
 as $v \to 0$ or $v \to \infty$
 $t^{-1+1/\theta}E(t,f) \to 0$ as $v \to 0$ or $v \to \infty$

(see [1, Thm 7.1.7]). As a result, we get

$$\int_0^\infty \left(v^{-\vartheta} K_\infty(v,a) \right)^q \frac{dv}{v} = -\frac{1}{\vartheta q} \int_0^\infty K_\infty(v,a)^q dv^{-\vartheta q} = \frac{1}{\vartheta q} \int_0^\infty v^{-\vartheta q} dK_\infty(v,a)^q.$$

On the other hand, integrating by parts with the change v = t/E(t, a), we get for $s + 1 = 1/\vartheta$ that

$$\int_0^\infty \left(v^{-\vartheta} K_\infty(v,a) \right)^q \frac{dv}{v} = \frac{1}{\vartheta q} \int_0^\infty \left(t/E(t,a) \right)^{-\vartheta q} dt^q = \frac{1}{\vartheta q^2} \int_0^\infty \left(t^s E(t,a) \right)^{\vartheta q} \frac{dt}{t}.$$

From the definition K_{∞} and K_2 , it follows that

$$K_{\infty}(t,a) \le K_2(t,a) \le 2^{1/2} K_{\infty}(t,a)$$
 (8)

(see [15, Remark 3.1]). Now, taking into account that

$$\frac{1}{\vartheta q^2}\|a\|_{\mathcal{B}^s_\tau}^{\vartheta q} = \frac{1}{\vartheta q^2} \int_0^\infty \left(t^s E(t,a)\right)^{\vartheta q} \frac{dt}{t} = \int_0^\infty \left(v^{-\vartheta} K_\infty(v,a)\right)^q \frac{dv}{v},$$

according to the left inequality from (8), we obtain

$$\frac{1}{\vartheta q^2} |a|_{\mathcal{B}^s_{\tau}}^{\vartheta q} \le \int_0^\infty \left(v^{-\vartheta} K_2(v, a) \right)^q \frac{dv}{v} = |a|_{(\mathfrak{A}_0, \mathfrak{A}_1)_{\vartheta, q}}^q. \tag{9}$$

By using the right inequality from (8), we have

$$\begin{aligned} |a|_{(\mathfrak{A}_0,\mathfrak{A}_1)_{\vartheta,q}}^q &= \int_0^\infty \left(v^{-\vartheta} K_2(v,a)\right)^q \frac{dv}{v} \leq 2^{q/2} \int_0^\infty \left(v^{-\vartheta} K_\infty(v,a)\right)^q \frac{dv}{v} \\ &= \frac{2^{q/2}}{\vartheta q^2} \int_0^\infty \left(t^s E(t,a)\right)^{\vartheta q} \frac{dt}{t} = \frac{2^{q/2}}{\vartheta q^2} |a|_{\mathcal{B}^q_\tau}^{\vartheta q} \end{aligned}$$

As a result, by virtue of (9) we have the inequalities

$$|a|_{(\mathfrak{A}_0,\mathfrak{A}_1)_{\vartheta,q}}^q \le \frac{2^{q/2}}{\vartheta q^2} |a|_{\mathcal{B}^s_{\tau}}^{\vartheta q} \le 2^{q/2} |a|_{(\mathfrak{A}_0,\mathfrak{A}_1)_{\vartheta,q}}^q \quad \text{with} \quad \tau = \vartheta q. \tag{10}$$

That gives the following isomorphism with equivalent quasinorms

$$\mathcal{B}_{\tau}^{s}(\mathfrak{A}_{0},\mathfrak{A}_{1}) = (\mathfrak{A}_{0},\mathfrak{A}_{1})_{\vartheta,q}^{1/\vartheta}.$$
 (11)

By [1, Lemma 7.1.2 and Thm 7.1.1] for each v > 0 there is t > 0 such that

$$(t^s E(t,a))^{\vartheta} \le v^{-\vartheta} K_{\infty}(v,a) \le (t^s E(t-0,a))^{\vartheta}. \tag{12}$$

Since $|a|_{\mathcal{B}_{\infty}^s}^{\vartheta} \leq |a|_{(\mathfrak{A}_0,\mathfrak{A}_1)_{\vartheta,q}}$ for $q=\infty$, the inequalities (12) for any v>0 yield

$$v^{-\vartheta} K_2(v, a) \le v^{-\vartheta} 2^{1/2} K_{\infty}(v, a) \le 2^{1/2} \left(t^s E(t - 0, a) \right)^{\vartheta}$$
$$\le 2^{1/2} \left(\sup_{t > 0} t^s E(t, a) \right)^{\vartheta} = 2^{1/2} |a|_{\mathcal{B}_{\infty}^s}^{\vartheta}.$$

As a result, $|a|_{(\mathfrak{A}_0,\mathfrak{A}_1)_{\vartheta,q}} \leq 2^{1/2}|a|_{\mathcal{B}_{\infty}^s}^{\vartheta}$ and the isomorphism (11) holds for $q=\infty$. Prove the Bernstein inequality (28). Combining (7) and (9), we obtain

$$|a|_{\mathcal{B}^{s}_{\tau}}^{\vartheta} \leq \begin{cases} \left[\frac{q}{1-\vartheta}\right]^{1/q} |a|_{0}^{1-\vartheta}|a|_{1}^{\vartheta} & \text{if } q < \infty\\ |a|_{0}^{1-\vartheta}|a|_{1}^{\vartheta} & \text{if } q = \infty. \end{cases}$$

$$(13)$$

Thus, the second inequality (28) we get by setting $s + 1 = 1/\vartheta$ and $\tau = \vartheta q$. Prove the inequality (27). Integrating $\min(1, v/t)K_2(t, a) \leq K_2(v, a)$, we obtain

$$\int_0^\infty \left(v^{-\vartheta}\min(1,v/t)\right)^q \frac{dv}{v} K_2(t,a)^q \leq \int_0^\infty \left(v^{-\vartheta} K_2(v,a)\right)^q \frac{dv}{v} = |a|_{(\mathfrak{A}_0,\mathfrak{A}_1)_{\vartheta,q}}^q.$$

Since the left integral can be rewritten as the following sum

$$\int_0^\infty \left(v^{-\vartheta}\min(1,v/t)\right)^q \frac{dv}{v} = \int_0^t v^{(1-\vartheta)q-1}t^{-q}dv + \int_t^\infty v^{-\vartheta q-1}dv = \frac{1}{t^\vartheta N_\vartheta^{1/q}},$$

we obtain the inequality

$$\bigg(\int_0^t \frac{v^{(1-\vartheta)q-1}}{t^q} dv + \int_t^\infty v^{-\vartheta q-1} dv\bigg)^{1/q} K_2(t,a) = \frac{K_2(t,a)}{t^\vartheta N_{\vartheta,q}} \leq |x|_{(\mathfrak{A}_0,\mathfrak{A}_1)_{\vartheta,q}}.$$

As a result, $K_2(t,a) \leq t^{\vartheta} N_{\vartheta,q} |a|_{(\mathfrak{A}_0,\mathfrak{A}_1)_{\vartheta,q}}$. Hence, taking into account (8) and (12), we get

$$v^{1-\vartheta}E(v,a)^{\vartheta} \le t^{-\vartheta}K_{\infty}(t,a) \le N_{\vartheta,q}|a|_{(\mathfrak{A}_0,\mathfrak{A}_1)_{\vartheta,q}}.$$

Applying (10), we obtain

$$v^{1-\vartheta}E(v,a)^{\vartheta} \le 2^{1/2}N_{\vartheta,a}|a|_{\mathcal{B}^s}^{\vartheta}.$$

Taking into account that $\vartheta = 1/(s+1) = q/\tau$, we obtain the inequality (27) for all $0 < q < \infty$.

In the case $q = \infty$, we have

$$t^s E(t,a) \le \sup_{t>0} t^s E(t,a) = |a|_{\mathcal{B}^s_\infty}$$

for all $a \in \mathcal{B}_{\infty}^{s}$. Thus, the inequality (27) holds for both cases.

Finally note that Bernstein-Jackson inequalities are achieved at $\tau = \infty$, so they are sharp.

Corollary 2 If the quasi-normed Abelian groups \mathfrak{A}_0 , \mathfrak{A}_1 are compatible then the following isomorphism with equivalent quasi-norms holds,

$$\mathcal{B}_{\tau}^{s}(\mathfrak{A}_{0},\mathfrak{A}_{1}) = (\mathfrak{A}_{0},\mathfrak{A}_{1})_{\vartheta,q}^{1/\vartheta}, \quad s+1 = 1/\vartheta, \quad \tau = \vartheta q.$$
 (14)

If compatible groups \mathfrak{A}_0 and \mathfrak{A}_1 are complete, the approximation group $\mathcal{B}_{\tau}^s(\mathfrak{A}_0,\mathfrak{A}_1)$ is complete.

Proof The isomorphism (14) immediately follows from (11) and Theorem 1. Note that if both compatible groups \mathfrak{A}_0 and \mathfrak{A}_1 are complete, the interpolation group $(\mathfrak{A}_0,\mathfrak{A}_1)_{\vartheta,q}$ is complete [1, Thm 3.4.2 & Lemma 3.10.2]. By Theorem 1 $\mathcal{B}^s_{\tau}(\mathfrak{A}_0,\mathfrak{A}_1)$ is complete.

Corollary 3 Let $(\mathfrak{A}_0, \mathfrak{A}_1)$, $(\mathfrak{A}'_0, \mathfrak{A}'_1)$ be compatible. If $T: \mathfrak{A}_i \to \mathfrak{A}'_i$ (i = 0, 1) are quasi-Abelian mappings, i.e. $T(a_0 + a_1) = b_0 + b_1$ and $|Ta_i|_{\mathfrak{A}_i} \leq \kappa_i |b_i|_{\mathfrak{A}'_i}$ with constants κ_i [1, p.81], then there exists a constant C > 0 such that

$$T \colon \mathcal{B}^s_\tau(\mathfrak{A}_0,\mathfrak{A}_1) \to \mathcal{B}^s_\tau(\mathfrak{A}_0',\mathfrak{A}_1') \quad and \quad |Ta|_{\mathcal{B}^s_\tau(\mathfrak{A}_0',\mathfrak{A}_1')} \leq C|a|_{\mathcal{B}^s_\tau(\mathfrak{A}_0,\mathfrak{A}_1)}.$$

Proof It follows from (10), (11) and the boundness of

$$T \colon (\mathfrak{A}_0, \mathfrak{A}_1)_{\vartheta, q} \to (\mathfrak{A}'_0, \mathfrak{A}'_1)_{\vartheta, q},$$

where $C \leq 4^{1/\vartheta} (\vartheta q^2)^{-1/\vartheta q} \kappa_0^s \kappa_1$ that gives the required inequality.

Corollary 4 Let $0 \le \vartheta_0 < \vartheta_1 \le 1$, $0 < q \le \infty$ and $\vartheta = (1 - \eta)\vartheta_0 + \eta\vartheta_1$ $(0 < \eta < 1)$. If the quasi-normed Abelian groups $(\mathfrak{A}_0, \mathfrak{A}_1)$, $(\mathfrak{A}'_0, \mathfrak{A}'_1)$ are compatible and the inclusions

$$(\mathfrak{A}_0,\mathfrak{A}_1)_{\vartheta_i,q}\subset \mathfrak{A}_i'\subset (\mathfrak{A}_0,\mathfrak{A}_1)_{\vartheta_i,\infty},\quad (i=0,1)$$

are valid, the following isomorphism with equivalent quasinorms holds,

$$\mathcal{B}_{\vartheta q}^{(1-\vartheta)/\vartheta}(\mathfrak{A}_0,\mathfrak{A}_1) = \mathcal{B}_{\eta q}^{(1-\vartheta)/\vartheta}(\mathfrak{A}_0',\mathfrak{A}_1'). \tag{15}$$

8

Proof Using (11) and the known equality $(\mathfrak{A}_0, \mathfrak{A}_1)_{\vartheta,q} = (\mathfrak{A}'_0, \mathfrak{A}'_1)_{n,q}$ [15], [8, Theorem 3] in the case r=1, we get the reiteration identity (15) with accuracy to quasinorm equivalency.

Corollary 5 The following continuous embedding hold

$$\mathcal{B}_{\vartheta q}^{(1-\vartheta)/\vartheta}(\mathfrak{A}_0,\mathfrak{A}_1) \hookrightarrow \mathcal{B}_{\vartheta p}^{(1-\vartheta)/\vartheta}(\mathfrak{A}_0,\mathfrak{A}_1) \quad \textit{with} \quad q < p.$$

Proof It follows from (11), since for arbitrary q < p the following continuous embedding $(\mathfrak{A}_0,\mathfrak{A}_1)_{\vartheta,q} \hookrightarrow (\mathfrak{A}_0,\mathfrak{A}_1)_{\vartheta,p}$ is true by virtue of [8, Lemma p. 385].

Corollary 6 If
$$\vartheta_0 < \vartheta_1$$
 then the embedding $\mathfrak{A}_1 \subset \mathfrak{A}_0$ yields
$$\mathcal{B}_{\vartheta_1 q}^{(1-\vartheta_1)/\vartheta_1}(\mathfrak{A}_0,\mathfrak{A}_1) \subset \mathcal{B}_{\vartheta_0 q}^{(1-\vartheta_0)/\vartheta_0}(\mathfrak{A}_0,\mathfrak{A}_1).$$

Proof It follows from Corollary 4 and [1, Theorem 3.4.1(d)].

Exact constants in bilateral error estimates

Let $(\mathfrak{A}, |\cdot|)$ be a κ -normed Abelian group. Consider a quasi-normed space $L^p_\mu = L^p_\mu(\mathfrak{A})$ with a positive Radon measure μ on a measure space (X,μ) of measurable functions $f: X \to \mathfrak{A}$ endowed with the κ -norm

$$||f||_p = \begin{cases} \left(\int |f(x)|^p \, \mu(dx) \right)^{1/p} & \text{if } 0$$

where supp $f \subset X$ is a measurable subset such that $f|_{X \setminus \text{supp } f} = 0$ and $f \neq 0$ almost everywhere on supp f with respect to the measure μ .

For a μ -measurable function $f: X \to \mathfrak{A}$, we define the distribution function

$$m(\sigma, f) = \mu \left\{ x \in X \colon |f(x)| > \sigma \right\}.$$

Denote by f^* the decreasing rearrangement of f, where

$$f^*(t) := \inf \{ \sigma > 0 \colon m(\sigma, f_\sigma) \le t \}$$

with $f_{\sigma}(x) = f(x)$ if $|f(x)| > \sigma$ and $f_{\sigma}(x) = 0$ otherwise (for details see [1, no 1.3]). The decreasing rearrangement f^* is nonnegative nonincreasing continuous on the right function of σ on $(0,\infty)$ which is equimeasurable with f in the sense that

$$m(\sigma, f) = m(\sigma, f^*)$$
 for all $\sigma \ge 0$.

In other words, the decreasing rearrangement of a function f is a generalized inverse of its distribution function in the sense that if $m(\sigma, f)$ is one-to-one then f^* is simply the inverse of $m(\sigma, f)$.

Since f is rearrangeable, $\{\sigma > 0 : m(\sigma, f_{\sigma}) \le t\}$ is nonempty for t > 0 thus f^* is finite on its domain.

In the case where $\mu(X) < \infty$, we consider f^* as a function on $(0, \mu(X))$, since $f^*(t) = 0$ for all $t > \mu(X)$. The set $\{\sigma > 0 : m(\sigma, f_\sigma) = 0\}$ can be empty. If f is bounded then $f^* \to ||f||_{\infty}$ as $t \to 0_+$, otherwise, f^* is unbounded at the origin.

As is also know (see e.g. [13]) for $f \in L^p_\mu$ with $p \ge 1$ that

$$\int |f|^p \, \mu(dx) = p \int_0^\infty \sigma^{p-1} m(\sigma, f) \, d\sigma, \quad \|f\|_p^p = \int_0^\infty f^*(t)^p \, dt.$$

The couple $(L_{\mu}^{0}, L_{\mu}^{\infty})$ is compatible in the interpolation theory sense (see e.g. [1, Lemma 3.10.3] or [8, no 1]). Given elements $f = f_{0} + f_{\infty}$ in the algebraic sum $L_{\mu}^{0} + L_{\mu}^{\infty}$, we define the best approximation *E*-functional $E(t, f; L_{\mu}^{0}, L_{\mu}^{\infty})$ with t > 0 as (see e.g. [1, no 7])

$$E(t,f) = E(t,f; L_{\mu}^{0}, L_{\mu}^{\infty}) = \inf \{ \|f - f_{0}\|_{\infty} \colon \|f_{0}\|_{0} < t \}, \quad f \in L_{\mu}^{\infty}.$$

In what follows, we use the equivalent parameters $0 < \vartheta < 1, \ 0 < q \le \infty$, where $s+1=1/\vartheta$ and $\tau=\vartheta q$. Following [10], the appropriate system of exact constants $C_{\theta,q}$ is defined by the normalization factor

$$N_{\theta,q} = \left(\int_0^\infty |t^{-\theta} \mathcal{N}(t)|^q \frac{dt}{t}\right)^{-1/q} \quad \text{with} \quad \mathcal{N}(t) = \frac{t}{\sqrt{1+t^2}}$$

in the known interpolation Lions-Peetre method to be

$$C_{\theta,q} = \begin{cases} \frac{2^{1/2\theta}}{(q^2\theta)^{1/q\theta}} N_{\theta,q}^{1/\theta} & \text{if } q < \infty, \\ \left(\frac{\sin \pi\theta}{\pi\theta}\right)^{1/2\theta} & \text{if } q = 2, \\ 2^{1/2\theta} & \text{if } q = \infty. \end{cases}$$

$$(16)$$

Remark 1 For this equivalent parameter system the sharp constant $c_{s,\tau}$ takes an equivalent form

$$2^{-1/2\theta}C_{\theta,q} = \begin{cases} c_{s,\tau} & \text{if } \tau < \infty \\ 1 & \text{if } \tau = \infty \end{cases},$$

where the following dependencies between indexes are performed,

$$2^{-1/2\theta}C_{\theta,q} = [(1-\theta)/q]^{1/q\theta} = (\theta q^2)^{-1/q\theta}N_{\theta,q}^{1/\vartheta}.$$




Fig. 1 3D-graph of $c_{s,\tau} = (s/\tau)^{1/\tau} (s+1)^{-2/\tau}$ for $s,\tau > 0$

A graphical image of the normalized constant sharp constant $c_{s,\tau}$ in variables s,τ shown on Fig. 1.

Theorem 7 (a) For any $0 < \vartheta < 1$ and $0 < q \le \infty$ the equality

$$f^*(t) = E(t, f; L^0_\mu, L^\infty_\mu), \quad f \in L^\infty_\mu$$
 (17)

and, as a consequence, the following equalities hold,

$$||f||_{q\theta^{2}/(1-\theta)} = \begin{cases} \left(\int_{0}^{\infty} \left[t^{-1+1/\theta} f^{*}(t) \right]^{q\theta} \frac{dt}{t} \right)^{1/q\theta} & \text{if } q < \infty \\ \sup_{0 < t < \infty} t^{-1+1/\theta} f^{*}(t) & \text{if } q = \infty. \end{cases}$$
(18)

(b) The Jackson-type inequality

$$f^*(t) \le t^{1-1/\theta} C_{\theta,q} \|f\|_{q\theta^2/(1-\theta)} \quad \text{for all} \quad f \in L^{q\theta^2/(1-\theta)}_{\mu},$$
 (19)

as well as, the following bilateral Bernstein-Jackson-type inequalities are valid,

$$t^{-1+1/\theta} f^*(t) \le C_{\theta,q} \|f\|_{q\vartheta^2/(1-\theta)} \le 2^{1/2\theta} \|f\|_0^{-1+1/\theta} \|f\|_{\infty}$$
 (20)

for all $f \in L^0_\mu \cap L^\infty_\mu$.

Proof (a) By definition, the functional E(t, f) is the infimum of $||f - f_0||_{\infty}$ such that $\mu(\text{supp } f_0) \leq t$. The next reasoning extends [1, Lemma 7.2.1] on the case of \mathfrak{A} -valued functions.

Put $g_0(x) = f(x)$ on the subset supp f and let $g_0(x) = 0$ outside supp f. Then we obtain $||f - g_0||_{\infty} \le ||f - g||_{\infty}$.

In a similar way, let $g_{\sigma}(x) = f(x)$ if $|f(x)| > \sigma$ and $g_{\sigma}(x) = 0$ otherwise. Then for the number $\tau = \sup\{|f(x)| \colon x \neq \text{supp } f\}$ we get $\sup f_{\tau} \subset \text{supp } f$. It follows $\mu(\sup g_{\tau}) \leq t$. Since $\|f - g_{\tau}\|_{\infty} \leq \tau$ and $\|f - g_{0}\|_{\infty} = \tau$, we obtain

$$E(t, f; L_{\mu}^{0}, L_{\mu}^{\infty}) = \inf_{\sigma} \left\{ \|f - g_{\sigma}\|_{\infty} \colon \mu \left(\operatorname{supp} g_{\sigma} \right) \le t \right\},\,$$

where the rigth hand side is equal to f^* . As a result, the equality (17) holds.

Using the best approximation E-functional, we define the quasi-normed space

$$E_{\theta,q}(L^0_{\mu}, L^\infty_{\mu}) = \left\{ f \in L^0_{\mu} + L^\infty_{\mu} : ||f||_{E_{\theta,q}} < \infty \right\},\tag{21}$$

$$||f||_{E_{\theta,q}} = \begin{cases} \left(\int_0^\infty \left[t^{-1+1/\theta} E(t,f) \right]^{q\theta} \frac{dt}{t} \right)^{1/q\theta} & \text{if } q < \infty, \\ \sup_{0 < t < \infty} t^{-1+1/\theta} E(t,f) & \text{if } q = \infty. \end{cases}$$
 (22)

By the known approximation theorem (see [1, Thm 7.2.2]) the isometric isomorphism

$$E_{\theta,q}(L_{\mu}^{0}, L_{\mu}^{\infty}) = L_{\mu}^{q\theta^{2}/(1-\theta)}, \quad \|f\|_{E_{\theta,q}} = \|f\|_{\vartheta^{2}q/(1-\theta)}$$
 (23)

holds for all $f \in L^{q\theta^2/(1-\theta)}_{\mu}$. Combining equalities (17), (21) and (23), we get the equality (18).

(b) In what follows, we use the classic integral of the real interpolation method

$$||f||_{\theta,q} = \left(\int_0^\infty \left| t^{-\theta} f(x) \right|^q \frac{dt}{t} \right)^{1/q}, \quad 0 < \theta < 1, \ 0 < q < \infty.$$
 (24)

Let us consider the quadratic K_2 -functional (see e.g. [12, App. B]) for the interpolation couple of quasi-normed groups $\left(L^0, L^\infty\right)$,

$$\begin{split} K_2(t,f) &= K_2 \left(t,f; \ L_{\mu}^0, L_{\mu}^{\infty} \right) \\ &= \inf_{f = f_0 + f_{\infty}} \left\{ \left(\|f_0\|_0^2 + t^2 \|f_{\infty}\|_{\infty}^2 \right)^{1/2} : f_0 \in L_{\mu}^0, f_{\infty} \in L_{\mu}^{\infty} \right\}, \end{split}$$

determining the real interpolation quasi-normed Abelian group

$$(L_{\mu}^{0}, L_{\mu}^{\infty})_{\theta, q} := K_{\theta, q} (L_{\mu}^{0}, L_{\mu}^{\infty}) = \{ f = f_{0} + f_{\infty} \colon ||K_{2}(\cdot, f)||_{\theta, q} < \infty \}$$

which is endowed with the norm

$$||f||_{K_{\theta,q}} = \begin{cases} N_{\theta,q} ||K_2(\cdot,f)||_{\theta,q} & \text{if } q < \infty \\ \sup_{t \in (0,\infty)} t^{-\theta} K_2(t,f) & \text{if } q = \infty. \end{cases}$$

Note that for any $0 < \vartheta < 1$ and q = 2 the following equalities hold,

$$N_{\theta,2} = \|\mathcal{N}\|_{\theta,2}^{-1} = \|K_2(\cdot,1)\|_{\theta,2}^{-1} = \left(\frac{2\sin\pi\theta}{\pi}\right)^{1/2}.$$

In fact, the relationship between the weight function $\mathcal{N}(t)^2 = t^2/(1+t^2)$ and the quadratic K_2 -functional is explained by the formula

$$N_{\theta,2} = \|\mathcal{N}\|_{\theta,2}^{-1} = \|K_2(\cdot,1)\|_{\theta,2}^{-1}$$

(see e.g. [12, Ex. B.4]). It follows from

$$\min_{z=z_0+z_1} \left(\alpha_0 |z_0|^2 + \alpha_1 |z_1|^2 \right) = \alpha_0 \alpha_1 |z|^2 \alpha_0 + \alpha_1$$

for a fixed $\alpha_0, \alpha_1 > 0$ and a complex z. This minimum is achieved when

$$\alpha_0 z_0 = \alpha_1 z_1 = \frac{\alpha_0 \alpha_1 z}{(\alpha_0 + \alpha_1)}.$$

Thus, $K_2(t,1)$ is minimized when f_0 , f_1 are such that

$$f_0 = t^2 f_1 = \frac{t^2}{1 + t^2}.$$

By integrating the above functions (see e.g. [12, Ex. B.5, Thm B.7]) it follows that the normalization factor $N_{\theta,2}$ is equal to $N_{\theta,2}=(2\sin\pi\theta/\pi)^{1/2}$.

In particular, the equalities (17),

$$f^*(t) = \inf \{ \sigma \colon m(\sigma, f_\sigma) \le t \}$$

$$=\inf\{\|f-f_0\|_{\infty}\colon \|f_0\|_0 < t\} = E(t,f;\ L_{\mu}^0,L_{\mu}^\infty), \quad f \in L_{\mu}^\infty,$$

as well as, the isometric isomorphism (23),

$$E_{\theta,q}(L^0_\mu, L^\infty_\mu) = L^{q\theta^2/(1-\theta)}_\mu, \quad \|f\|_{E_{\theta,q}} = \|f\|_{\vartheta^2q/(1-\theta)}$$

allows calculating the exact form of best constants (16).

Corollary 8 The decreasing rearrangement for q = 2 has the estimation

$$f^*(t) \le t^{1-1/\theta} \left(\frac{\sin \pi \theta}{\pi \theta}\right)^{1/2\theta} ||f||_{2\theta^2/(1-\theta)} \quad \text{for all} \quad f \in L_\mu^{2\theta^2/(1-\theta)}$$
 (25)

which for $\theta = 1/2$ can be written as follows

$$f^*(t) \le \frac{2}{\pi t} \int |f| \mu(dx) \quad \text{for all} \quad f \in L^1_\mu.$$
 (26)

Note that if $\theta \to 1$ then $\sin \pi \theta \to 0$ thus above inequalities disappear. One has a sense only for the quasi-normed group $L_{\mu}^{2\theta^2/(1-\theta)}$ with $0 < \theta < 1$.

Corollary 9 The equality (18) with $s + 1 = 1/\vartheta$ and $\tau = \vartheta q$ takes the form

$$||f||_{\tau/s} = \begin{cases} \left(\int_0^\infty \left[t^s f^*(t) \right]^\tau \frac{dt}{t} \right)^{1/\tau} & \text{if } \tau < \infty \\ \sup_{0 < t < \infty} t^s f^*(t) & \text{if } \tau = \infty \end{cases}$$

for any $f \in L^{\tau/s}_{\mu}$. Then the Jackson-type inequality takes the form

$$f^*(t) \le t^{-s} 2^{(s+1)/2} c_{s,\tau} ||f||_{\tau/s} \quad \text{for all} \quad f \in L_{\mu}^{\tau/s}$$
 (27)

and bilateral Bernstein-Jackson-type inequalities take the form

$$t^{s} 2^{-(s+1)/2} f^{*}(t) \le c_{s,\tau} \|f\|_{\tau/s} \le \|f\|_{0}^{s} \|f\|_{\infty} \quad \text{for all} \quad f \in L_{\mu}^{0} \cap L_{\mu}^{\infty}. \tag{28}$$

It instantly follows from the relations

$$C_{\theta,q} = \begin{cases} 2^{(s+1)/2} c_{s,\tau} & \text{if } \tau < \infty \\ 2^{(s+1)/2} & \text{if } \tau = \infty \end{cases}, \quad \text{where} \quad \vartheta = \frac{1}{s+1}, \quad 2^{1/2\theta} = 2^{(s+1)/2}.$$

Remark 2 The inequalities (28) is an extension on the case measurable functions the known best approximation Bernstein-Jackson inequalities. Whereas the inequality (27) is an extension of the approximation Jackson inequality. The scale of approximation quasi-normed Abelian groups is often denoted as

$$\mathcal{B}_{\tau}^{s}(L_{\mu}^{0}, L_{\mu}^{\infty}) := E_{\theta, q}(L_{\mu}^{0}, L_{\mu}^{\infty}),$$

where the space \mathcal{B}_{τ}^{s} coincides with a suitable extension of the Besov type quasinormed groups (see e.g. [1, Thm 7.2.4], [18, Def. 4.2.1/1]). The scale of quasi-normed Besov Abelian groups for another approximation couples is described in [7].

4 Examples of bilateral estimates with exact constants

Taking into account the previous statements, we can give typical examples of Bernstein-Jackson inequalities with explicit constants for different types of best approximations.

Example 1 (Applications to classic Besov scales) Let $\mathfrak{A}_0 = L^p(\mathbb{R}^n)$ with $(1 . As is known (see e.g. [14]) each real-valued function <math>a \in L^p(\mathbb{R}^n)$ can be approximated by entire analytic functions $g \in \mathcal{E}_p^t(\mathbb{C}^n)$, where $\mathcal{E}_p^t(\mathbb{C}^n)$ means the space of entire analytic functions on \mathbb{C}^n of an exponential type t > 0 with restrictions to \mathbb{R}^n belonging to $L^p(\mathbb{R}^n)$. The best approximations can be characterized by the best approximation functional

$$E\left(t, a; \, \mathcal{E}_p, L^p(\mathbb{R}^n)\right) = \inf_{t>0} \left\{ \|a - g\|_{L^p(\mathbb{R}^n)} \colon g \in \mathcal{E}_p^t(\mathbb{C}^n) \right\},\,$$

where the subspace $\mathfrak{A}_1 = \mathcal{E}_p$ with $\mathcal{E}_p = \bigcup_{t>0} \mathcal{E}_p^t(\mathbb{C}^n)$ is endowed with the quasinorm

$$|g|_{\mathcal{E}_p} = ||g||_{L^p(\mathbb{R}^n)} + \{\sup |\zeta| \colon \zeta \in \operatorname{supp} \hat{g}\}, \quad g \in \mathcal{E}_p$$

defined using the support supp \hat{g} of the Fourier-image \hat{g} . Then according to Theorem 1 the corresponding approximation inequalities take the form

$$t^{s}E\left(t, a; \ \mathcal{E}_{p}, L^{p}(\mathbb{R}^{n})\right) \leq c_{s,\tau} \|f\|_{B^{s}_{p,\tau}(\mathbb{R}^{n})} \leq 2^{1/2} |a|_{\mathcal{E}_{p}}^{s} \|a\|_{L^{p}}, \ a \in \mathcal{E}_{p} \cap L^{p}(\mathbb{R}^{n}),$$
$$E\left(t, a; \ \mathcal{E}_{p}, L^{p}(\mathbb{R}^{n})\right) \leq t^{-s} c_{s,\tau} \|a\|_{B^{s}_{p,\tau}(\mathbb{R}^{n})}, \quad a \in B^{s}_{p,\tau}(\mathbb{R}^{n}),$$

In this case the approximation scale

$$\mathcal{B}_{\tau}^{s}(\mathcal{E}_{p}, L^{p}(\mathbb{R}^{n})) = (\mathcal{E}_{p}, L^{p}(\mathbb{R}^{n}))_{\mathfrak{H}, q}^{1/\vartheta}$$

exactly coincides with the classic Besov scale denoted by $B_{p,\tau}^s(\mathbb{R}^n)$ (see [18, p.197]).

Example 2 (Applications to scales of periodic functions) Following e.g. [1, no 1.5], [17] we can write Bernstein-Jackson inequalities in a more general form. Let $X=\mathbb{T}$ be the 1-dimensional torus and the Hilbert space $\mathfrak{A}_1=L^2(\mathbb{T})$ has the orthonormal basis $\left\{\mathfrak{e}_k=e^{2\pi i kt}\colon k\in\mathbb{Z},\,t\in\mathbb{T}\right\}$. Let $\mathfrak{A}_0=\{\text{trigonometric polynomials }a_0\}$ with the quasi-norm $|a_0|_0=\deg a_0$, and $\mathfrak{A}_1=\{2\pi\text{-periodic functions }a\}$ with the norm $|a|_1=\|a\|_{L^2(\mathbb{T})}$, as well as, $E(n,a;\,\mathfrak{A}_0,\mathfrak{A}_1)=\inf\left\{\|a-a_0\|_{L^2(\mathbb{T})}\colon\deg a_0< n\right\}$ for

all
$$a \in L^2(\mathbb{T})$$
. Let $c_{j,\tau} := \left\{ \begin{bmatrix} \frac{j}{\tau(j+1)^2} \end{bmatrix}^{1/\tau} & \text{if } \tau < \infty \\ 1 & \text{if } \tau = \infty \end{bmatrix} \right.$ with $j \in \mathbb{N}$. Then Bernstein-

Jackson inequalities have the form

$$n^{j}E(n,a) \leq c_{j,\tau}|a|_{\mathcal{B}^{j}_{\tau}} \leq 2^{1/2}|a|_{0}^{j}|a|_{1} \quad \text{for all} \quad a \in \mathfrak{A}_{0} \cap \mathfrak{A}_{1},$$

$$E(n,a) \leq n^{-j}c_{j,\tau}|a|_{\mathcal{B}^{j}_{\tau}} \quad \text{for all} \quad a \in \mathcal{B}^{j}_{\tau}(\mathfrak{A}_{0},\mathfrak{A}_{1}),$$

where
$$\mathcal{B}_{\tau}^{s}(\mathfrak{A}_{0},\mathfrak{A}_{1})=(\mathfrak{A}_{0},\mathfrak{A}_{1})_{\vartheta,q}^{1/\vartheta}$$
 with $j+1=1/\vartheta\in\mathbb{N},\,\tau=\vartheta q.$

Let $\mathfrak{A}=L^2(\mathbb{T})$. Consider the Banach space L^p_μ of $L^2(\mathbb{T})$ -valued functions $f\colon X\to L^2(\mathbb{T})$. Then bilateral Bernstein-Jackson inequalities have the form

$$n^{-j}2^{-(j+1)/2}f^*(n) \le c_{j,\tau}||f||_{\tau/j} \le ||f||_0^j||f||_{\infty}, \quad f \in L^0_\mu \cap L^\infty_\mu,$$

$$f^*(n) \le n^j 2^{(j+1)/2} c_{j,\tau} ||f||_{\tau/j}, \quad f \in L_{\mu}^{\tau/j},$$

where $f^*(n) = E(n, f; L^0_\mu, L^\infty_\mu) = \inf \{ \sigma \colon m(\sigma, f_\sigma) \le t \}$ for all $f \in L^\infty_\mu$.

Example 3 (Applications to operator spectral approximations) Consider the example commonly used in foundations of quantum systems for describing quantum states (see e.g. [9, no 3.4]). Similar estimates of spectral approximations were early analyzed in the paper [7]. In what follows, we consider the spaces $L^p = L^p(\mathbb{R}; \mathbb{R})$ of real-valued functions.

Let H be a Hilbert complex space with the norm $\|\cdot\|_H = \langle\cdot|\cdot\rangle^{1/2}$ and a self-adjoint unbounded linear operator T with the dense domain $\mathcal{D}(T) \subset H$ is given. By spectral theorem the measurable function f from T can be well defined using the following spectral expansions (see e.g. [9])

$$T = \int_{\sigma(T)} \lambda \, \mu(d\lambda), \quad f(T) = \int_{\sigma(T)} f(\lambda) \, \mu(d\lambda),$$

where μ is a unique projection-valued measure determined on its spectrum $\sigma(T) \subset \mathbb{R}$ with values in the Banach space of bounded linear operators $\mathcal{L}(H)$, which can be extended on $\mathbb{R} \setminus \sigma(A)$ as zero.

At beginning, we consider the case when for any Borel set $\Sigma \subset \mathbb{R}$ and any $\psi \in \mathcal{D}(A)$ the $\mathcal{L}(H)$ -valued measurable function

$$\Sigma \longmapsto f(\Sigma) = \int_{\Sigma} f(\lambda) \, \mu_{\psi}(d\lambda)$$

belongs to L^{∞} with respect to the positive measure $\mu_{\psi}(d\lambda) := \langle \mu(d\lambda)\psi \mid \psi \rangle$, where $\|\psi\|_{H} = 1$.

Consider the corresponding quadratic forms $f_{\psi}(\Sigma) = \langle f(\Sigma)\psi | \psi \rangle$ as measurable functions of L^2 and let f_{ψ}^* be the suitable decreasing rearrangement. By Theorem 11 the inequalities (27) and (28) in this case take the form

$$f_{\psi}^{*}(t) \le t^{1-1/\theta} C_{\theta,q} \|f_{\psi}\|_{q\theta^{2}/(1-\theta)}, \quad f_{\psi} \in L_{\mu}^{q\theta^{2}/(1-\theta)},$$
 (29)

$$t^{-1+1/\theta} f_{\psi}^{*}(t) \leq C_{\theta,q} \|f_{\psi}\|_{q\vartheta^{2}/(1-\theta)} \leq 2^{1/2\theta} \|f_{\psi}\|_{0}^{-1+1/\theta} \|f_{\psi}\|_{\infty}, \quad f_{\psi} \in L_{\mu}^{0} \cap L_{\mu}^{\infty},$$
(30)

where $||f_{\psi}||_{\infty} = \sup_{\Sigma \subset \mathbb{R}} |f_{\psi}(\Sigma)|$ and $||f_{\psi}||_{0} = \mu_{\psi}$ (supp f_{ψ}). Applying the estimation (26) from Corollary 8, we obtain

$$f_{\psi}^*(t) \le \frac{2}{\pi t} \int |f(\lambda)| \, \mu(d\lambda) \quad \text{for all} \quad f_{\psi} \in L_{\mu}^1.$$
 (31)

The inequality (29), (31) give error estimations (in quadratic forms terms) of spectral approximations $f_{\psi}(T) = \langle f(T)\psi \mid \psi \rangle$ by $f_{\psi}(\Sigma)$ with $\mu(\Sigma) \leq t$. The inequalities (30) characterise the approximation accuracy.

In another case, using the $\mathcal{L}(H)$ -valued measurable uniformly bounded functions

$$\tilde{f} \colon \Sigma \longmapsto \int_{\Sigma} f(\lambda) \, \mu(d\lambda), \quad f \in L^1_{\mu}$$

belonging to $L^1_{\mu}(\mathbb{R},\mathcal{L}(H))$, we same as above obtain the inequality

$$\tilde{f}^*(t) \le \frac{2}{\pi t} \left\| \int f(\lambda) \, \mu(d\lambda) \right\|_{\mathcal{L}(H)}.$$




Fig. 2 Graphs of the function $f(t) = 10\sqrt{\frac{2}{\pi t^3}}e^{-\frac{(t-2)^2}{2t}}$ (dots) and its non-increasing rearrangement $f^*(t)$ (polygon), (0 < t < 10)




Fig. 3 Graphs of approximation E(u, f) via $u^{-s}c_{s,\tau}||f||_{\mathcal{B}^{s}_{\tau}}$ $(0 < u < 11, s = \tau = 2)$ using Jackson's inequality

Example 4 (Samples of numerical calculations) The inequality (27) together with the exact value of the normalized constant $c_{s,\tau}$ are suitable for direct numerical calculations. Illustrate this on several graphs, taking as an example the inverse Gaussian distribution function.

Consider two-parameter inverse Gaussian distribution with support on $(0, \infty)$ that is given by

$$f(t) = C\sqrt{\frac{l}{2\pi t^3}} \exp\left(-\frac{l(t-m)^2}{2m^2t}\right), \quad C, t > 0$$

with the mean m > 0 and the shape parameter l > 0. By Theorem 11

$$f^*(u) = E(u, f) = \inf \{ ||f - f_{\sigma}||_{\infty} \colon m(\sigma, f_{\sigma}) \le u \},$$

$$m(\sigma, f_{\sigma}) = \mu \{ u \in (0, \infty) \colon |f_{\sigma}(u)| > \sigma \}$$

coincides with the non-increasing rearrangement of f on $(0, \infty)$. As a result, we obtain

$$f^*(u) = E(u, f) \le c_{s,\tau} u^{-s} ||f||_{\tau/s}$$
 for all $f \in L_{\mu}^{\tau/s}$,

where $L_{\mu}^{\tau/s} = \mathcal{B}_{\tau}^{s} \left(L_{\mu}^{0}, L_{\mu}^{\infty} \right)$ in accordance with Remark 2 and the equality (18). The obtained results of numerical calculations are illustrated on Fig. 2,3,4.

Statements and Declarations: There are no conflicts and potential competing of interest to disclose.

References

- [1] Bergh, J., Löfström, J.: Interpolation Spaces. Springer, Berlin-Göttingen-Heidelbergm (1976)
- [2] Bernd, C.: Inequalities of Bernstein-Jackson-type and the degree of compactness of operators in Banach spaces. Ann. Inst. Fourier (Grenoble). 35(3) 79–118 (1985)
- [3] Burchard, A., Hajaiejb, H.: Rearrangement inequalities for functionals with monotone integrands. J. Funct. Anal. 233. 561–582 (2006).
- [4] Grafakos, L.: Classical Fourier analysis, vol. 249/250 Springer (2014)
- [5] DeVore, R.A., Popov, V.: Interpolation of Besov spaces, Trans. Amer. Math. Soc. 305, 397–414 (1988)
- [6] DeVore, R.A.: Nonlinear approximation. Acta Numer. 7(1), 51–150 (1998)
- [7] Dmytryshyn, M., Lopushansky, O.: On spectral approximations of unbounded operators. Complex Anal. Oper. Theory. 13(8), 3659–3673 (2019)
- [8] Komatsu, N.: A general interpolation theorem of Marcinkiewicz type. Tôhoku Math. Journ., 33, 383–393 (1981)
- [9] Hall, B. C.: Quantum Theory for Mathematicians. Springer (2013)
- [10] Lopushansky, O.: Bernstein–Jackson Inequalities on Gaussian Hilbert Spaces. J. Fourier Anal. Appl., 29, 57–78 (2023)
- [11] Maligranda, L., Persson, L.E.: The E-functional for some pairs groups. Results Math. 20(1/2), , 538-553 (1991)
- [12] McLean, W.: Strongly Elliptic Systems and Boundary Integral Equations. Cambridge Univ. Press (2000)
- [13] O'Neil, R., Weiss, O.: The Hilbert transform and rearraangement of functions, Stud. Math., 23, 190–197 (1963)
- [14] Nikolskii, S.: Approximation of functions of several variables and imbedding theorems. Springer, Berlin-Göttingen-Heidelberg (1975)
- [15] Peetre, J., Sparr, G.: Interpolation of normed Abelian groups. Ann. Mat. Pura Appl. 92(1), 217–262 (1972)

- [16] Pietsch, A.: Approximation spaces. J. Approx. Theory 32(2), 115–134 (1981)
- [17] Prestin, J., Savchuk, V.V., Shidlich, A.L.: Direct and inverse theorems on the approximation of 2- periodic functions by Taylor Abel Poisson operators, Ukr. Math. J. 69(5), 766–781 (2017)
- [18] Triebel, H.: Interpolation Theory. Function Spaces. Differential Operators. North-Holland Publ. (1978)