# The Lamperti transformation in the infinite-dimensional setting and the genealogies of self-similar Markov processes

Arno Siri-Jégousse<sup>1</sup> and Alejandro H. Wences<sup>2</sup>

<sup>1</sup>IIMAS, Universidad Nacional Autónoma de México, México <sup>2</sup>CIMI-LAAS, Université de Toulouse, CNRS, Toulouse, France

May 28, 2024

### Abstract

We propose a change in focus from the prevalent paradigm based on the branching property as a tool to analyze the structure of population models, to one based on the self-similarity property, which we also introduce for the first time in the setting of measure-valued processes. By extending the well-known Lamperti transformation for self-similar Markov processes to the Banach-valued case we are able to generalize celebrated results in population genetics that describe the frequency-process of measure-valued stable branching processes in terms of the subfamily of Beta-Fleming-Viot processes. In our work we describe the frequency process of populations whose total size evolves as any positive self-similar Markov process in terms of general  $\Lambda$ -Fleming-Viot processes. Our results demonstrate the potential power of the self-similar perspective for the study of population models in which the reproduction dynamics of the individuals depend on the total population size, allowing for more complex and realistic models.

Keywords: self-similarity, Lamperti transformation, measure-valued processes, coalescent processes.

MSC 2020: 92D25, 60G18, 60J68.

### 1 Introduction and Main Results

In the same spirit as in [5, 32], the work of [8] describes the genealogy of a population that evolves as an stable measure-valued branching process, say  $(\mu_t)_{t\geq 0}$ , indexed by a parameter  $\beta \in (0, 2]$ . Formally, the latter are Markov processes on the state space  $\mathbb{M}([0, 1])$  of finite positive measures on the type space [0, 1], and with generator  $\mathcal{F}$ having one of the two forms:

1) 
$$\mathcal{F}F(\mu) = \int_{\mathcal{T}} \mu(da) \int_0^\infty h^{-1-\beta} dh \left\{ F(\mu + h\delta_a) - F(\mu) - hF'(\mu; a) \right\}, \text{ for } \beta \in (0, 2),$$
  
or

### 1 INTRODUCTION AND MAIN RESULTS

2)  $\mathcal{F}F(\mu) = \int_{\mathcal{T}} \mu(da) F''(\mu; a, a)$ , which heuristically corresponds to the case  $\beta = 2$ .

In the latter, we have removed the drift terms and the constants for simplicity. The genealogy of such models is described by means of characterizing the associated frequency process  $\left(\frac{\mu t}{\|\mu_t\|}\right)_{t\geq 0}$ , where  $\|\mu\| := \mu([0,1])$ . The authors show that, after the time change  $c_{\beta-1}(t) = \inf\{s \geq 0: \int_0^s \|\mu_u\|^{1-\beta} du > t\}$ , the frequency process becomes Markov in itself and is a member of the Beta subfamily of  $\Lambda$ -Fleming-Viot (FV) processes [4]. The genealogy of the underlying population is then understood via the well-know duality relation between  $\Lambda$ -FV processes and  $\Lambda$ -coalescents [4].

The family of  $\Lambda$ -FV (resp.  $\Lambda$ -coalescents) generalize the standard FV process (resp. the Kingman coalescent) to capture the population genetics dynamics of a wider range of neutral populations, including populations with highly skewed offspring distributions (but also many more models, including models with selection). Still, [8] proved that only the Beta subfamily can be obtained from a branching process by using their method based on path-wise random time changes. For the rest of the superprocesses (measure-valued branching processes), the associated frequency process is not Markov in itself under any random time change that is written in terms of the total population size (see their Lemma 3.5 for further details). This apparent lack of a branching process counterpart for the rest of the Lambda family has motivated research seeking variants of the main results in [8] to obtain different  $\Lambda$ -FV (resp.  $\Lambda$ -coalescents) for the frequency process (resp. genealogy) of branching processes, but using different transformations. An example is the culling procedure in [10] (see also [20]) who work with the two-dimensional counterparts of FV processes, and who approximate populations with constant size and in distribution, losing the path-wise quality. Another example is the work of [25] who study the genealogy of multi-type branching processes locally in time by computing the corresponding coalescence rates, which notably depend on the size of the populations of each of the finitely-many types. In our case it has motivated a change of perspective by now focusing on the self-similarity property instead, enabling us to use robust path-wise tools such as the Lamperti transformation in the infinite-type setting. This technique also allows us to incorporate dependencies in the reproduction dynamics of the individuals, mainly through the total population size, and also to accommodate in a simple way the effect of this size on the coalescence rates and the genealogy.

A measure-valued Markov process  $(\mu_t)_{t\geq 0}$ , and more generally a Banach-valued Markov process  $(X_t)_{t\geq 0}$ , is said to satisfy the <u>self-similarity (SS)</u> property with index  $\alpha \in \mathbb{R}$  (denoted  $\alpha$ -SS) if

(1) 
$$\forall a > 0, \quad \{X_t, \mathbb{P}_x^X\} = \{aX_{a^{-\alpha}t}, \mathbb{P}_{a^{-1}x}^X\}.$$

On the other hand, a two-coordinate process  $(Z_t)_{t\geq 0} = (\rho_t, \xi_t)_{t\geq 0}$  on  $S \times \mathbb{R}$ , with S being the unit sphere of a Banach space, and with lifetime  $\zeta^Z$ , is said to be a Markov additive process (MAP) if for any  $y \in S, z \in \mathbb{R}, s, t \geq 0$ , and for any positive measurable function f on  $S \times \mathbb{R}$ , one has

(2) 
$$\mathbb{E}_{\theta,z}^{Z}\left[f(\rho_{t+s},\xi_{t+s}-\xi_{t}),t+s<\zeta^{Z}|\mathcal{F}_{t}\right]=\mathbb{E}_{\rho_{t},0}^{Z}\left[f(\rho_{s},\xi_{s}),s<\zeta^{Z}\right]\mathbb{1}_{t<\zeta^{Z}}.$$

We obtain the following (also see Theorem 4.2 below for further details).

### 1 INTRODUCTION AND MAIN RESULTS

**Theorem 1.1.** There exists a standard measure-valued  $\alpha$ -SS Markov process  $(\mu_t)_{t\geq 0}$  with generator of the form

$$\mathcal{F}F(\mu) = \frac{1}{\|\mu\|^{\alpha}} \left( \mathcal{G}_{\kappa}^{(D)}F(\mu) + \mathcal{G}_{\sigma}^{(B)}F(\mu) + \mathcal{G}_{\Lambda}^{(J)}F(\mu) \right)$$

where, for  $\sigma \geq 0, \kappa \in \mathbb{R}$ , and  $\Lambda \in M((0,1))$ , the operators are defined by

$$\begin{aligned} \mathcal{G}_{\kappa}^{(D)}F(\mu) &= \int_{\mathcal{T}} \mu(da)\kappa F'(\mu;a), \\ \mathcal{G}_{\sigma}^{(B)}F(\mu) &= \|\mu\| \int_{\mathcal{T}} \mu(da) \frac{\sigma^2}{2} F''(\mu;a,a), \text{ and} \\ \mathcal{G}_{\Lambda}^{(J)}F(\mu) &= \int_{\mathcal{T}} \frac{\mu(da)}{\|\mu\|} \int_{(0,1)} \frac{\Lambda(d\zeta)}{\zeta^2} \left\{ F\left(\mu + \|\mu\| \frac{\zeta}{1-\zeta} \delta_a\right) \\ &- F(\mu) - \|\mu\| \left(|\log(1-\zeta)| \,\mathbb{1}_{\zeta < 1/2}\right) F'(\mu;a) \right\} \end{aligned}$$

The total size of the population  $(\|\mu_t\|)_{t\geq 0}$  is a general positive  $\alpha$ -SS Markov process with non-negative jumps. Furthermore, letting

$$c_{\alpha}(t) = \inf\{s \ge 0: \int_0^s \|\mu_u\|^{-\alpha} \, du > t\}$$

the processes  $\left(\frac{\mu_{c(t)}}{\|\mu_{c(t)}\|}, \log(\|\mu_{c(t)}\|)\right)_{t\geq 0}$  is a MAP. Its first coordinate is a  $(\Lambda + \frac{\sigma^2}{2}\delta_0)$ -Fleming-Viot process; whereas the second coordinate is a spectrally-positive Lévy process.

Through an appropriate choice of the parameters  $\sigma$ ,  $\kappa$ ,  $\Lambda$ , the above theorem recovers the cases when the stability index  $\beta \in (1, 2)$  of [8], which correspond to measure-valued branching processes that also enjoy the  $(\beta - 1)$ -SS property, as seen through an adaptation of the result in [28] from  $\mathbb{R}$ -valued to measure-valued processes.

Our method is based on a generalization of the Lamperti transformation [30, 1, 11] to the infinite-dimensional setting, which we now describe (see section 2 for further details). Let E be a conic subset of a normed vector space  $(\mathbb{V}, \|\cdot\|)$ , and suppose that  $(E, \mathbf{d})$  is a metric space such that the map  $x \to \|x\|$  is continuous. In the application above, E is the space of positive measures  $\mathbb{M}(\mathcal{T})$  over a type space  $\mathcal{T}$ , and  $\mathbf{d}$  is a metrization of the weak topology on  $\mathbb{M}(\mathcal{T})$ . We also assume  $(E, \mathbf{d})$  to be locally-compact and second-countable, and augment it to  $\hat{E} = E \cup \{\infty\}$  where  $\infty$  is a point at infinity if E is not compact, or just an isolated point if E is compact.

We consider only standard Markov process with values in  $D([0, \infty), \hat{E})$ . The latter is the space of càdlàg functions with values in  $\hat{E}$  that are absorbed at  $\{\infty, 0\}$ . Also, for a general process X we write  $\zeta_{\infty}^X$  for its absorption time to  $\infty$ ,  $\zeta_0^X$  for its absorption time to  $0 \in E$ , and define its lifetime  $\zeta^X := \zeta_{\infty}^X \wedge \zeta_0^X$ . The self-similar Lamperti transformation can be expressed as follows.

**Theorem 1.2.** Let  $(X_t)_{t\geq 0}$  be a standard  $\alpha$ -SS Markov process. Consider the additive functional  $t \to \int_0^t ||X_u||^{-\alpha} du$  for  $t \in [0, \zeta^X]$ , and its generalized inverse

$$c_{\alpha}(t) := \inf \left\{ s > 0 \colon \int_{0}^{s} \|X_{u}\|^{-\alpha} \, du \ge t \right\}, \quad t \in [0, \int_{0}^{\zeta^{X}} \|X_{u}\|^{-\alpha} \, du].$$

Then the process  $(Y_t)_{t\geq 0} = \left(\frac{X_{c_{\alpha}(t)}}{\|X_{c_{\alpha}(t)}\|}, \log(\|X_{c_{\alpha}(t)}\|)\right)_{t\geq 0}$  is a standard MAP with lifetime  $\zeta^Y = \zeta^Y_{\infty} \wedge \zeta^Y_0 \underset{a.s.}{=} \int_0^{\zeta^X} \|X_u\|^{-\alpha} du.$ 

Conversely, let  $(\rho_t, \xi_t)_{t\geq 0}$  be a standard MAP, and let  $\zeta^{\xi} = \zeta_{\infty}^{\xi} \wedge \zeta_0^{\xi}$  be its lifetime, after which it is absorbed at some extra state. For any  $\alpha \geq 0$ , consider the inverse additive functional

$$\gamma_{\alpha}(t) \coloneqq \inf \left\{ s > 0 \colon \int_{0}^{s} e^{\alpha \xi_{u}} du \ge t \right\}, \quad t \in [0, \int_{0}^{\zeta^{\xi}} e^{\alpha \xi_{u}} du].$$

Then the process  $(X_t)_{t\geq 0} = \left(\rho_{\gamma_{\alpha}(t)}e^{\xi_{\gamma_{\alpha}(t)}}\right)_{t\geq 0}$  is a standard  $\alpha$ -SS Markov process with lifetime  $\zeta^X = \zeta^X_{\infty} \wedge \zeta^S_0 \underset{a.s.}{\overset{\zeta^{\xi_{\infty}} \wedge \zeta^{\xi}_0}{\overset{\zeta^{\xi_{\infty}}}}{\overset{\zeta^{\xi_{\infty}}}{\overset{\zeta^{\xi_{\infty}}}}{\overset{\zeta^{\xi_{\infty}}}{\overset{\zeta^{\xi_{\infty}}}{\overset{\zeta^{\xi_{\infty}}}}{\overset{\zeta^{\xi_{\infty}}}{\overset{\zeta^{\xi_{\infty}}}{\overset{\zeta^{\xi_{\infty}}}}{\overset{\zeta^{\xi_{\infty}}}{\overset{\zeta^{\xi_{\infty}}}{\overset{\zeta^{\xi_{\infty}}}}}}}}}}}}}}}}$ 

Our results show an unexplored link between the fields of mathematical population genetics and self-similar Markov processes in infinite dimensions, motivating new research and opening new questions in both of these fields separately, but also at their intersection. For instance, the populations driven by  $\mathcal{F}$  above, characterized by four parameters (the Lévy triplet and the self-similarity index) constitute only a sub-class of the entire family of measure-valued self-similar processes which is yet to be fully described. Whether one will need a new model to describe their corresponding genealogies, outside or extending the family of  $\Lambda$  (and more generally  $\Xi$ ) coalescent processes, is still an open question. At the same time, measure-valued processes, together with the well-established analytic tools available in population genetics such as duality methods, may serve as a suitable template for the development of the theory of self-similar Markov processes and their Lamperti transformations in infinite dimensions.

The rest of the manuscript is structured as follows. Section 2 is devoted to the development of our main methodological result, the Lamperti transformation in the infinite-dimensional setting (Theorem 1.2). On the other hand, our main phenomenological result (Theorem 1.1) is developed in section 4, which is preceded by section 3 in which we describe some necessary preliminaries on coalescent processes, FV processes, and Dawson-Watanabe processes. In turn, section 4 is decomposed into three main subsections, section 4.1 providing details and expanding upon Theorem 1.1, and sections 4.2 and 4.3 which constitute the two main steps in the construction of an "intermediary" process ( $\nu_t$ )<sub>t≥0</sub> with generator of the form  $\mathcal{G} = \mathcal{G}_{\kappa}^{(D)} + \mathcal{G}_{\sigma}^{(B)} + \mathcal{G}_{\Lambda}^{(J)}$  from which the process ( $\mu_t$ )<sub>t≥0</sub> can be obtained via a Lamperti time change.

### 2 Lamperti Transformations in Normed Spaces

In this section we get inspiration from the works of [30, 1, 11] and generalize the selfsimilar Lamperti transformation to processes taking values in a conic subset E of a normed space  $(\mathbb{V}, \|\cdot\|)$ . In particular E is closed under non-negative scalar transformations (i.e.  $x \in E$  implies  $ax \in E$  for all  $a \ge 0$ ), and we can furthermore define a polar decomposition from E to  $S \times \mathbb{R}$ , where S is the intersection of E with the unit sphere in  $\mathbb{V}$ , as  $x \mapsto (x/||x||, ||x||)$ . In fact all we need are these two properties for E and that the function  $\|\cdot\|$  satisfies ||ax|| = |a| ||x||, but we subscribe to the normed space setting in order to avoid technicalities. We also recall that we have assumed the metric space (E, d) to be locally-compact and second-countable, and that  $\hat{E}$  is its one-point compactification if E is not compact, and  $\infty$  is an isolated point if E is compact. Furthermore, we also assume that the function  $\|\cdot\|$  is continuous on (E, d).

Most of the arguments used in our main proofs can already be found in [1, 22], we expand them to general state spaces and also deal with possible explosion of the processes involved.

We assume that all the processes that we consider are standard in the sense of Definition 9.2 in [9]; namely that they satisfy the following:

- i) their respective filtrations  $(\mathcal{F}_t)_{t>0}$  are right-continuous,
- ii) they are absorbed at  $\infty$  at time  $\zeta_{\infty} \in [0, \infty]$ ,
- iii) they are càdlàg and quasi-left-continuous on  $[0, \zeta_{\infty})$ ,
- iv) they are strong-Markov processes with measurable probability kernels  $P_t(x, \cdot)$  from  $\hat{E}$  to  $\hat{E}$  corresponding to the law of the process at time t.

Let us call  $D([0,\infty), \hat{E})$  the trajectory space of such processes, i.e. the space of trajectories on  $\hat{E}$  that are càdlàg on  $[0, \zeta_{\infty})$  and have constant value  $\infty$  on  $[\zeta_{\infty}, \infty)$ . Endow  $D([0,\infty), \hat{E}) \subset \hat{E}^{\mathbb{R}_+}$  with  $\mathcal{F}$ , the trace  $\sigma$ -algebra induced by  $\sigma(\pi_t; t \geq 0)$  where  $\pi_t$  is the projection at time t. Note that when E is a locally-compact secondcountable metric space, the extended space  $\hat{E}$  becomes compact and metrizable (see e.g. Proposition VII.1.15 im [29]) and  $\mathcal{F}$  coincides with the Borel  $\sigma$ -algebra induced by the Skorohod topology on  $D([0,\infty), \hat{E})$  (Theorem 12.5 in [6]). Also we will denote by  $\mathbb{P}_x(\cdot)$  the law of the processes on  $\hat{E}^{\mathbb{R}_+}$ , started at x. Finally  $\zeta_x$  will denote the reaching time of a process to an absorbing state x (which could be an extra state, or the point at infinity  $\infty$ ); whereas  $\zeta$  will denote the lifetime of the process. Typically  $\zeta = \zeta_{\infty} \wedge \zeta_0$ . We will often decorate  $\zeta, \mathcal{F}$ , and  $\mathbb{P}$  with superscripts specifying the distinct processes we will deal with, as below.

An  $\hat{E}$ -valued Markov process  $(X_t)_{t>0}$  that is absorbed at  $\{\infty, 0\}$  and has lifetime  $\zeta^X = \zeta_0^X \wedge \zeta_\infty^X$  is said to satisfy the <u>self-similarity (SS)</u> property with index  $\alpha \in \mathbb{R}$  ( $\alpha$ -SS) if (1) holds. Alternatively, it is easily seen that the process X is  $\alpha$ -SS if and only if for all  $t \geq 0, a > 0, x \in E, A \in \mathcal{B}(E)$ , its transition kernels  $P_t^X$  satisfy

$$P_t^X(x,A) = P_{a^{-\alpha}t}^X \left( a^{-1}x, a^{-1}A \right).$$

Secondly, an  $\hat{E}$ -valued Markov process Y absorbed at  $\{\infty, 0\}$  with lifetime  $\zeta^Y = \zeta_0^Y \wedge \zeta_\infty^Y$  is said to be scalar multiplicative homogeneous (SMH) if its transition kernels  $P_t^Y$  satisfy, for all  $t \ge 0, u > 0, x \in E, A \in \mathcal{B}(E)$ ,

(3) 
$$P_t^Y(x,A) = P_t^Y(ux,uA),$$

or, in other words, if

$$\forall u > 0, \{Y_t, \mathbb{P}_x^Y\} = \{uY_t, \mathbb{P}_{u^{-1}x}^Y\}.$$

Thirdly, recalling that S is the intersection of E with the unit sphere in V, a  $S \times \mathbb{R}$ -valued Markov process  $(Z_t)_{t\geq 0} = (\rho_t, \xi_t)_{t\geq 0}$  that has lifetime  $\zeta^Z = \zeta_{\infty}^{\xi}$  (the explosion time of  $\xi$  on the one-point compactification of  $\mathbb{R}$ , after which Z is absorbed at some extra state), is said to be a Markov additive process (MAP) if (2) holds. Equivalently, it is easily seen that the process  $(\rho_t, \xi_t)_{t\geq 0}$  is a MAP if and only if, for all  $t \geq 0, a \in \mathbb{R}, (\theta, z) \in S \times \mathbb{R}^+$  and  $A \in \mathcal{B}(S \times \mathbb{R}^+)$ ,

(4) 
$$P_t^{Z}((\theta, z), A) = P_t^{Z}((\theta, z+a), A+(0, a));$$

where we have written  $A + (0, a) := \{(\rho, z + a) : (\rho, z) \in A\}$ . This can be interpreted as saying that Z is additive-homogeneous on the second coordinate.

We will establish transformations between SS and SMH processes on the one hand, and between SMH processes and MAPs on the other. In fact, the latter is simply a bijection given by the "log-polar decomposition" isomorphism  $\Phi: E \setminus \{0\} \to S \times \mathbb{R}$ defined as  $\Phi(x) = (x/||x||, \log(||x||))$ . We have the following.

**Proposition 2.1** (SMH  $\iff$  MAP). Let Y be a SMH Markov process with trajectories in  $D([0,\infty), \hat{E})$  and absorbed at 0, and set  $\zeta^Y = \zeta_0^Y \wedge \zeta_\infty^Y$ . Then  $(Z_t)_{t\geq 0} = (\Phi(Y_t))_{t\geq 0}$  is a MAP with lifetime  $\zeta^Z = \zeta^Y$ .

Conversely, if  $(Z_t)_{t\geq 0} = (\rho_t, \xi_t)_{t\geq 0}$  is a MAP with lifetime  $\zeta^Z = \zeta_{\infty}^{\xi}$ , then  $Y = (\Phi^{-1}(Z_t))_{t\geq 0} \equiv (e^{\xi_t}\rho_t)_{t\geq 0}$  is a SMH Markov process with lifetime  $\zeta^Y = \zeta^Z$ .

*Proof.* Given that  $\Phi$  is bijective and continuous except at the absorbing state 0, it is clear that the transformed processes are standard whenever the starting process is. Thus we need only verify (4) in the first case, and (3) in the second. We only do this for the first. We have, assuming (3) in the second equality below, and for all  $t \geq 0, a \in \mathbb{R}, (\theta, z) \in S \times \mathbb{R}, A \in \mathcal{B}(S \times \mathbb{R}),$ 

$$P_t^Z((\theta, z), A) = P_t^Y(e^{z\theta}, \Phi^{-1}(A)) = P_t^Y(e^{z+a\theta}, e^{a\Phi^{-1}(A)})$$
$$= P_t^Z((\theta, z+a), A + (0, a)).$$

The transformation between SS and SMH processes is given in terms of random time changes. The proof follows the heuristics in the proof of Theorem 2.3 in [1] but adapted to our setting.

**Theorem 2.2** (Self-Similar Lamperti Time Change). Let  $(X_t)_{t\geq 0}$  be a standard Markov process with trajectories in  $D([0,\infty), \hat{E})$ . Let  $\zeta_{\infty}^X$  be its explosion time,  $\zeta_0^X$  be its absorption time to  $0 \in E$ , and define its lifetime  $\zeta^X = \zeta_{\infty}^X \wedge \zeta_0^X$ . Consider the additive functional  $t \to \int_0^t ||X_u||^{-\alpha} du$  for  $t \in [0, \zeta^X]$ , and its generalized inverse

$$c_{\alpha}(t) := \inf \left\{ s > 0 \colon \int_{0}^{s} \|X_{u}\|^{-\alpha} du \ge t \right\}, \quad t \in [0, \int_{0}^{\zeta^{X}} \|X_{u}\|^{-\alpha} du]$$

- i) If  $(X_t)_{t\geq 0}$  is  $\alpha$ -SS, then the process  $(Y_t)_{t\geq 0} = (X_{c_{\alpha}(t)})_{t\geq 0}$  is a standard SMH Markov process with lifetime  $\zeta^Y = \zeta^Y_{\infty} \wedge \zeta^Y_0 = \int_0^{\zeta^X} ||X_u||^{-\alpha} du.$
- ii) Setting  $X_{\infty} = \infty$  if  $\alpha > 0$ , and  $X_{\infty} = 0$  if  $\alpha \le 0$ , the process  $(Y_t)_{t \ge 0}$  is the unique solution to

(5) 
$$a.s. \quad \forall t \ge 0, \quad Y_t = X_{\int_0^t ||Y_u||^\alpha du \wedge \zeta^X}.$$

Conversely, let  $(Y_t)_{t\geq 0}$  be a standard Markov process with trajectories in  $D([0,\infty), \hat{E})$ . Let  $\zeta^Y = \zeta^Y_{\infty} \wedge \zeta^Y_0$  be its lifetime, after which it is absorbed at some extra state. Consider, for any  $\alpha \geq 0$ , the inverse additive functional

$$\gamma_{\alpha}(t) := \inf \left\{ s > 0 \colon \int_{0}^{s} \|Y_{u}\|^{\alpha} \, du \ge t \right\}, \quad t \in [0, \int_{0}^{\zeta^{Y}} \|Y_{u}\|^{\alpha} \, du].$$

- iii) If  $(Y_t)_{t\geq 0}$  is SMH, then the process  $(X_t)_{t\geq 0} = (Y_{\gamma_{\alpha}(t)})_{t\geq 0}$  is a standard  $\alpha$ -SS Markov process with lifetime  $\zeta^X = \zeta^X_{\infty} \wedge \zeta^X_0 = \int_0^{\zeta^Y} ||Y_u||^{\alpha} du$ .
- iv) The process  $(X_t)_{t>0}$  is the unique solution to

(6) 
$$a.s. \ \forall t \ge 0, \quad Y_t = X_{\int_0^{t \land \zeta Y} \|Y_u\|^{\alpha} du},$$

that also satisfies  $X_t = X_{t \wedge \int_0^{\zeta^Y} ||Y_s||^{\alpha} ds}$ .

*Proof.* To ease notations, we will omit the index  $\alpha$  in  $c_{\alpha}(t)$  and  $\gamma_{\alpha}(t)$ .

By Propositions IV.1.6 and IV.1.13 in [9] (and the discussion around eq. IV.1.8 therein), the mappings  $t \to \int_0^t ||X_u||^{-\alpha} du$  and  $t \to \int_0^t ||Y_u||^{\alpha} du$  both define continuous strong additive functionals of  $((X_t)_{t\geq 0}, \zeta^X)$  and  $((Y_t)_{t\geq 0}, \zeta^Y)$  respectively (see Definitions IV.1.1 and IV.1.11 in [9]). Then, by Exercise V.2.11 iv) in [9], the time-changed processes  $(X_{c(t)})_{t\geq 0}$  and  $(Y_{\gamma(t)})_{t\geq 0}$  are strong Markov processes in each case. Since c(t) (resp.  $\gamma(t)$ ) is continuous on  $[0, \int_0^{\zeta^X} ||X_u||^{-\alpha} du)$  (resp.  $[0, \int_0^{\zeta^Y} ||Y_u||^{\alpha} du)$ ), it follows that  $(X_{c(t)})_{t\geq 0}$  (resp.  $(Y_{\gamma(t)})_{t\geq 0}$ ) is quasi-left continuous. Also, by Lemma 2.4 below, the mapping  $x \to \mathbb{E}_x^Y[f(Y_t)] \equiv \mathbb{E}_x^X[f(X_{c(t)})]$  is measurable for every  $t \geq 0$  and  $f \in \bar{\mathbb{B}}(\hat{E})$ , the space of bounded Borel functions on  $\hat{E}$ , so that the process defined by  $Y_t = X_{c(t)}$  in i)) is indeed a standard process; and similarly for the process defined by  $X_t = Y_{\gamma(t)}$  in iii)).

We now show that the process Y in i)) is SMH. Let  $\hat{c}(t)$  be the functional c(t) applied to the process  $(\hat{X}_t)_{t\geq 0} \coloneqq (aX_{a-\alpha t})_{t\geq 0}$ . Observe that the change of variable  $v = a^{-\alpha}u$  yields

$$a^{-\alpha}\hat{c}(t) = a^{-\alpha}\inf\left\{s \ge 0: \int_0^s \|aX_{a^{-\alpha}u}\|^{-\alpha} du \ge t\right\}$$
$$= \inf\left\{a^{-\alpha}s \ge 0: \int_0^{a^{-\alpha}s} \|X_v\|^{-\alpha} dv \ge t\right\}$$
$$= \inf\left\{s \ge 0: \int_0^s \|X_v\|^{-\alpha} dv \ge t\right\}$$
$$= c(t).$$

Thus  $\hat{X}_{\hat{c}(t)} = aX_{a^{-\alpha}\hat{c}(t)} = aX_{c(t)}$ . This, together wit the  $\alpha$ -SS property of X in the second equality below give, for  $A \in \mathcal{B}(\hat{E}), x \in \hat{E}$ , and a > 0,

$$\mathbb{P}_{x}^{Y}(Y_{t} \in A) = \mathbb{P}_{x}^{X}(X_{c(t)} \in A) = \mathbb{P}_{a^{-1}x}^{X}(\hat{X}_{\hat{c}(t)} \in A) = \mathbb{P}_{a^{-1}x}^{X}(aX_{c(t)} \in A)$$
$$= \mathbb{P}_{a^{-1}x}^{Y}(Y_{t} \in a^{-1}A).$$

We proceed similarly to prove iii). For a > 0 let  $\hat{\gamma}(t)$  be the functional  $\gamma(t)$  applied to the process  $(\hat{Y}_t)_{t>0} \coloneqq (aY_t)_{t\geq 0}$ . Observe that

$$\hat{\gamma}(t) = \inf\left\{s \ge 0: \int_0^s \|aY_u\|^\alpha \, du \ge t\right\}$$
$$= \inf\left\{s \ge 0: \int_0^s \|Y_u\|^\alpha \, du \ge a^{-\alpha}t\right\}$$
$$= \gamma(a^{-\alpha}t).$$

The above yields  $\hat{Y}_{\hat{\gamma}(t)} = aY_{\gamma(a^{-\alpha}t)}$ . This, together with the SMH property of Y in the second equality below give

$$\mathbb{P}_{x}^{X}(X_{t} \in A) = \mathbb{P}_{x}^{Y}(Y_{\gamma(t)} \in A) = \mathbb{P}_{a^{-1}x}^{Y}(\hat{Y}_{\hat{\gamma}(t)} \in A) = \mathbb{P}_{a^{-1}x}^{Y}(aY_{\gamma(a^{-\alpha}t)} \in A)$$
$$= \mathbb{P}_{a^{-1}x}^{X}(aX_{a^{-\alpha}t} \in A).$$

We now show ii). Write  $\psi(t) \coloneqq \int_0^t ||X_s||^{-\alpha} ds$ . If  $\alpha \ge 0$  note that, since  $(X_t)_{t\ge 0}$  is absorbed at  $\infty$ , we have  $\zeta^X = \zeta^X_{\infty}$  whenever  $\psi(\infty) < \infty$ , and  $\zeta^X_{\infty} = \infty$  whenever  $\psi(\infty) = \infty$ . In any case  $\psi$  is monotonic and absolutely continuous on comparison time  $\psi(\infty) = 0$ . intervals contained in  $[0, \zeta^X]$  with inverse function c(t) and a.e. derivative  $\|X_t\|^{-\alpha}$ . It follows that for any  $\psi(ds)$ -integrable function  $\beta_s$  on  $[0, \zeta^X]$ , we have

$$\int_{0}^{t} \beta_{s} \|X_{s}\|^{-\alpha} ds = \int_{0}^{\psi(t)} \beta_{c(s)} ds.$$

If  $t < \psi(\infty)$ , in particular if  $\psi(\infty) = \infty$ , we have  $0 < \|X_s\|^{\alpha} < \infty$  on  $s \in [0, c(t)]$  so that

$$c(t) = \int_0^{c(t)} \|X_s\|^{\alpha} \|X_s\|^{-\alpha} \, ds = \int_0^t \|X_{c(s)}\|^{\alpha} \, ds;$$

whereas if  $\infty > t \ge \psi(\infty)$  then

$$\int_0^t \left\| X_{c(s)} \right\|^{\alpha} ds \ge \lim_{u \uparrow \psi(\infty)} \int_0^u \left\| X_{c(s)} \right\|^{\alpha} ds = \lim_{u \uparrow \psi(\infty)} c(u) = c(\psi(\infty)) \equiv \zeta_{\infty}^X$$

In any case

$$c(t) \equiv c(t) \wedge \zeta^{X} = \int_{0}^{t} \|Y_{s}\|^{\alpha} \, ds \wedge \zeta^{X}.$$

In particular, recalling that we have set  $X_{\infty} = \infty$  if  $\alpha > 0$  and  $X_{\infty} = 0$  if  $\alpha \leq 0$ , we have

$$Y_t = X_{c(t)\wedge\zeta X} = X_{\int_0^t \left\| X_{c(s)} \right\|^\alpha ds \wedge \zeta X} = X_{\int_0^t \left\| Y_s \right\|^\alpha ds \wedge \zeta X}.$$

To see that this is the unique solution note that if  $Y'_t$  satisfies (5), then the function

$$\tilde{c}(t) = \int_0^t \left\| Y'_s \right\|^\alpha ds = \int_0^t \left\| X_{\tilde{c}(s)} \right\|^\alpha ds$$

is continuous, strictly increasing, and has derivative  $\|X_{\tilde{c}(s)}\|^{\alpha}$  a.e. on [0, t] whenever  $\tilde{c}(t) < \zeta^X$ . Then, in this case we have

$$t = \int_0^t \left\| X_{\tilde{c}(s)} \right\|^{\alpha} \left\| X_{\tilde{c}(s)} \right\|^{-\alpha} ds = \int_0^{\tilde{c}(t)} \left\| X_s \right\|^{-\alpha} ds$$

which implies  $\tilde{c}(t) = c(t)$  whenever  $\tilde{c}(t) < \zeta^X$  and  $\tilde{c}(\psi(\infty)) = \lim_{t \uparrow \psi(\infty)} c(t) = \zeta^X$ . Then since  $Y'_t = X_{\tilde{c}(t) \land \zeta^X}$ , we conclude  $Y'_t = X_{c(t) \land \zeta^X} = Y_t$  for all  $t \ge 0$ . Finally, we prove iv). Write  $\varphi(t) = \int_0^t ||Y_s||^{\alpha} ds$ . Note that  $\zeta^Y \le \gamma(\varphi(\infty))$  and  $\gamma(t \land s) = \gamma(t) \land \gamma(s)$  for every  $s, t \ge 0$ . Then, the fact that  $(X_t)_{t\ge 0}$  solves (6) follows from  $X_t \equiv Y_{\gamma(t \land \varphi(\zeta^Y) \land \varphi(\infty))}$  and the fact that  $\varphi$  is strictly increasing and continuous on  $[0, \zeta^Y]$  with left inverse  $\gamma$ . This yields

$$X_{\varphi(t\wedge\zeta^Y)} = Y_{\gamma\left(\varphi(t\wedge\zeta^Y)\wedge\varphi(\zeta^Y)\right)} = Y_{t\wedge\zeta^Y}.$$

The fact that  $(X_t)_{t\geq 0}$  is the unique solution follows from the fact that  $\varphi(\gamma(t)) = t$  on  $[0, \zeta^Y]$  which gives, for any other solution  $(X'_t)_{t\geq 0}$  to the functional equation,

$$Y_{t \wedge \zeta^Y} = X'(\varphi(\gamma(t) \wedge \zeta^Y)) = X'(t \wedge \varphi(\zeta^Y)).$$

The composition of the above two transformations between MAP and SMH processes, and between SMH and  $\alpha$ -SS processes respectively, leads to Theorem 1.2 which for  $\mathbb{R}^d$ -valued processes is the Lamperti transformation of [1]; and for  $\mathbb{R}^+$ -valued processes is the original transformation of [30]. We also refer the reader to the result in [11] and [21].

The following proposition gives a characterization of  $\alpha$ -SS and SMH processes in terms of their generators. For  $a \geq 0$ , let  $S_a$  be the operator that scales space by a factor of a, i.e. that takes  $f \in \overline{B}(E)$  to  $S_a f(x) = f(ax)$ .

**Proposition 2.3.** Let  $(X_t)_{t\geq 0}$  be a Markov process with generator having a core  $\mathcal{A} \subset \overline{B}(E) \times \overline{B}(E)$  such that  $S_a \overline{\mathcal{D}}(\mathcal{A}) \subset \mathcal{D}(\mathcal{A})$ . Then the following are equivalent.

- i) The process X is  $\alpha$ -SS.
- *ii)* For all  $a \geq 0$ ,  $\mathcal{A} = a^{-\alpha} \mathcal{S}_{a^{-1}} \mathcal{A} \mathcal{S}_a$ .

Similarly, let  $(Y_t)_{t\geq 0}$  be a Markov process with generator having a core  $\mathcal{A} \subset \overline{B}(E) \times \overline{B}(E)$  such that  $S_a \mathcal{D}(\mathcal{A}) \subset \mathcal{D}(\mathcal{A})$ . Then the following are equivalent.

- iii) The process Y is SMH.
- iv) For all  $a \geq 0$ ,  $\mathcal{A} = \mathcal{S}_{a^{-1}} \mathcal{A} \mathcal{S}_a$ .

*Proof.* We only prove equivalence between i) and ii), the proof of the SMH case being analogous. Assuming i) we obtain  $\mathbb{E}_x^X[f(X_t)] = \mathbb{E}_{a^{-1}x}^X[\mathcal{S}_a f(X_{a^{-\alpha}t})]$ , so that taking time derivatives,

$$\mathcal{A}f(x) = \frac{d}{dt}\Big|_{t=0} \mathbb{E}_x^X[f(X_t)]$$
  
=  $\frac{d}{dt}\Big|_{t=0} \mathbb{E}_{a^{-1}x}^X[\mathcal{S}_a f(X_{a^{-\alpha}t})]$   
=  $a^{-\alpha} \mathcal{A} \mathcal{S}_a f(a^{-1}x)$   
=  $a^{-\alpha} \mathcal{S}_{a^{-1}} \mathcal{A} \mathcal{S}_a f(x).$ 

Then, ii) follows. The converse implication follows from the fact that  $a^{-\alpha}S_{a^{-1}}\mathcal{A}S_a$  coincides with the generator of the Markov process  $\{aX_{a^{-\alpha}t}; \mathbb{P}_{a^{-1}x}^X, x \in \hat{E}\}$  on  $\mathcal{D}(\mathcal{A})$ . Then ii) implies that its semigroup coincides with that of  $\{X_t; \mathbb{P}_x^X, x \in \hat{E}\}$ , and the two processes are equal in distribution.

We end this section with the following technical lemma that was used in the proof of Theorem 2.2.

Lemma 2.4. We have the following measurability of mappings.

i) The mappings  $(x_t)_{t\geq 0} \to (c(t))_{t\geq 0}$  and  $(y_t)_{t\geq 0} \to (\gamma(t))_{t\geq 0}$  are measurable from  $D([0,\infty), \hat{E})$  to  $D^0([0,\infty), [0,\infty])$ , the non-decreasing elements of  $D([0,\infty), [0,\infty])$ , endowed with the relative  $\sigma$ -algebra inherited from the Skorohod  $\sigma$ -algebra in  $D([0,\infty), [0,\infty])$ .

ii) The mappings  $(x_t)_{t\geq 0} \to (x_{c(t)})_{t\geq 0}$  and  $(z_t)_{t\geq 0} \to (z_{\gamma(t)})_{t\geq 0}$  are measurable from  $D([0,\infty), \hat{E})$  to  $D([0,\infty), \hat{E})$ .

*Proof.* Recall that, E being locally compact and second countable, the Borel  $\sigma$ -algebra on  $D([0, \infty), \hat{E})$  generated by the Skorohod topology, and the trace  $\sigma$ -algebra generated by the finite-dimensional projections on  $D([0, \infty), \hat{E})$ , coincide. Then, i) follows from the fact that, for each  $t_0$ , the map  $(x_t)_{t\geq 0} \to c(t_0)$  is measurable from  $\mathcal{B}(D([0, \infty), \hat{E}))$  to  $\mathcal{B}(\hat{[}0, \infty])$ , where  $\mathcal{B}(\cdot)$  stands for the Borel  $\sigma$ -algebra in any topological space. Since  $\mathcal{B}(D([0, \infty), [0, \infty]))$  is generated by the finite dimensional projections (see section 12 in [6]), the latter implies that the map  $(x_t)_{t\geq 0} \to (c(t))_{t\geq 0}$  is measurable from  $D([0, \infty), \hat{E})$  to  $D([0, \infty), [0, \infty])$ .

On the other hand, ii) follows from i) plus Appendix M16 in [6].

## **3** Preliminary objects of study

In the following  $C(\cdot)$  (resp.  $B(\cdot)$ ) refers to the space of  $\mathbb{R}$ -valued continuous (resp. measurable) functions defined on some toplogical (resp. measurable) space; whereas  $\overline{C}(\cdot)$  (resp.  $\overline{B}(\cdot)$ ) refers to its bounded counterpart. Also  $C_0^k$  refers to the space of continuous functions vanishing at infinity, that have a continuous *k*-th derivative. Additionally,  $\mathcal{D}(\cdot)$  will refer to the domain of an operator.

### **3.1** $\Lambda$ -coalescents

We expose the construction of coalescents with multiple merger from the seminal works of [34, 35]. For a positive integer p, let  $[p] = \{1, \dots, p\}$  and  $\mathscr{P}_{[p]}$  be the space of partitions of [p] endowed with the discrete topology. We call the elements of any partition  $\pi \in \mathscr{P}_{[p]}$  the blocks of  $\pi$  and denote its number by  $\#\pi$ . Let  $\Lambda$  be a finite measure on [0, 1] which can be decomposed as

$$\Lambda = \Lambda(\{0\})\delta_0 + \mathbb{1}_{(0,1]}\Lambda.$$

The  $(p, \Lambda)$ -coalescent process  $(\Pi_t)_{t\geq 0}$  is a Markov jump process with values in  $\mathscr{P}_{[p]}$ that evolves through "coagulations" or mergers. The latter consists of constructing a new coarser partition of [p] from an initial  $\pi \in \mathscr{P}_{[p]}$  by taking the union of a collection of blocks that are present in  $\pi$ . The coagulations of  $(\Pi_t)_{t\geq 0}$  are directed by the measure  $\Lambda$  via the following rules; at time  $t \geq 0$ :

**Pairwise coagulations:** Any pair of blocks of  $\Pi_t$  coagulate at rate  $\Lambda(\{0\})$ .

**Coin-flip coagulations:** Any collection of  $2 \leq i \leq j = \#\Pi_t$  blocks of  $\Pi_t$ , coagulate into a single block at rate  $\beta_{j,i}^{(\Lambda)} \coloneqq \int_{(0,1]} \zeta^{i-2} (1-\zeta)^{j-i} \Lambda(d\zeta)$ .

The first dynamics correspond to those of Kingman's coalescent [27]. The second dynamics have the following well-known interpretation: at rate  $\zeta^{-2}\Lambda(d\zeta)$  a value  $\zeta \in (0,1]$  is drawn; then, each block of  $\Pi_t$  decides to participate in the coagulation event with probability  $\zeta$ . This representation for the rates implies that those processes are consistent according to p and can thus be extended to  $p = \infty$ . In this case, we will talk about  $\Lambda$ -coalescents.

A famous and important example of  $\Lambda$ -coalescent processes is the family of Beta coalescents [37, 19] in which  $\Lambda(d\zeta) = c\zeta^{1-\beta}(1-\zeta)^{\beta-1}d\zeta$  for  $\beta \in (0,2)$ . See [2] for a thorough exposition of general coalescent processes.

### **3.2** The $\Lambda$ -Fleming-Viot processes

We begin with a few remarks on the space  $M(\mathcal{T})$  endowed with the topology of weak convergence. By Theorem 1.14 in [31] the space  $M(\mathcal{T})$  is locally-compact whenever  $\mathcal{T}$  is compact. In fact, following the proof of this theorem, the set  $M_r(\mathcal{T}) = \{\mu \in M(\mathcal{T}) : \mu(\mathcal{T}) \leq r\}$ , for  $r \geq 0$ , is compact.

Let us write

$$\langle f, \mu \rangle \coloneqq \int \mu(da) f(a).$$

An important class of functions in  $C(M(\mathcal{T}))$  is the algebra of polynomials  $\mathtt{Pol}\,(M(\mathcal{T}))$  which is the linear span of monomials of the form

$$F_{\phi,p}(
ho) = \left\langle \phi, 
ho^{\otimes p} \right\rangle, \quad \phi \in \overline{\mathsf{C}}(\mathcal{T}^p).$$

By a straightforward extension of Lemma 2.1.2 in [12] (extending the arguments therein to  $\overline{C}(M(\mathcal{T}))$ ), the polynomials Pol  $(M(\mathcal{T}))$  are dense in the topology of uniform convergence on compact sets on  $C(M(\mathcal{T}))$ , and convergence determining for the topology of weak convergence in  $M(M_r(\mathcal{T}))$  for every  $r \geq 0$ .

A function  $F \in C(M(\mathcal{T}))$  is said to be differentiable if its derivative in the direction of  $a \in \mathcal{T}$  (more precisely of  $\delta_a$ ) given by

$$F'(\mu; a) \coloneqq \lim_{\epsilon \to 0} \frac{F(\mu + \epsilon \delta_a) - F(\mu)}{\epsilon},$$

exists and is continuous as a function of  $a \in \mathcal{T}$ . We denote by  $F''(\mu; a, b)$  the second derivative of F, first in the direction of a and then in the direction of b; whereas for higher derivatives we write  $F^{(\ell)}(\mu; a_1, \cdots, a_\ell)$  for the corresponding sequential derivatives in the directions of  $a_1, \cdots, a_\ell$ . By Lemma 2.1.2 in [12], the polynomials Pol  $(\mathfrak{M}(\mathcal{T}))$  are infinitely differentiable. Their derivatives are given by

$$F'_{\phi,p}(\mu;a) = \sum_{i=1}^{p} \left\langle \phi, \mu^{\otimes i-1} \otimes \delta_a \otimes \mu^{p-i-1} \right\rangle,$$

and recursively, for  $\boldsymbol{a} = (a_1, \cdots, a_\ell)$ ,

(7) 
$$F_{\phi,p}^{(\ell)}(\mu; \boldsymbol{a}) = \begin{cases} \sum_{\sigma \in \mathcal{P}(\boldsymbol{a}, \mu, p)} \langle \phi, \otimes_{i=1}^{p} \sigma_{i} \rangle & \text{if } \ell \leq p \\ 0 & \text{otherwise} \end{cases}$$

where the sum is over all the permutations  $\sigma \in \mathcal{P}(\boldsymbol{a}, \mu, p)$ , say  $\sigma = (\sigma_1, \dots, \sigma_p)$ , of the atomic measures  $\delta_{a_1}, \dots, \delta_{a_\ell}$  and  $p - \ell$  copies of  $\mu$ . Note that in particular

$$\left|F_{\phi,p}(\mu) + F'_{\phi,p}(\mu;a) + F''_{\phi,p}(\mu;a,b)\right| \le C \|\phi\| (1 + \dots + \|\mu\|^p)$$

for some C depending on p, so that the polynomials  $\mathtt{Pol}(\mathtt{M}(\mathcal{T}))$  are contained in the set

(8)  
$$D := \left\{ F \in C(M(\mathcal{T})) : \exists C > 0, p \ge 1 \text{ s.t.} \\ |F(\mu)| + |F'(\mu; a)| + |F''(\mu; a, a)| \le C(1 + \dots + |||\mu|||^p) \right\},$$

which will appear in the domain of generators of measure-valued processes further ahead.

Now, let  $\Lambda$  and  $\beta_{j,i}^{(\Lambda)}$  be as in section 3.1. The  $\Lambda$ -Fleming-Viot process [4] is the process with values in the space  $PM(\mathcal{T})$  of probability measures on  $\mathcal{T}$  and generator of the form

(9) 
$$\mathcal{R}F_{\phi,p}(\rho) = \int_{\mathcal{T}} \rho(da) \sum_{\ell=2}^{p} \beta_{p,\ell}^{(\Lambda)} \sum_{\sigma \in \mathcal{P}(a,\mu,p)} \left\{ \langle \phi, \otimes_{i=1}^{p} \sigma_{i} \rangle - \langle \phi, \rho^{\otimes p} \rangle \right\}$$
$$= \int_{\mathcal{T}} \rho(da) \sum_{\ell=2}^{p} \beta_{p,\ell}^{(\Lambda)} \left( F_{\phi,p}^{(\ell)}(\rho;a) - {p \choose \ell} F_{\phi,p}(\rho) \right);$$

where we have made a slight abuse of notation by writing  $F_{\phi,p}^{(\ell)}(\mu;a)$  for the  $\ell$ -times derivative of F, all in the direction of a. The above form of the generator yields the following well-known duality relation between  $\Lambda$ -Fleming-Viot processes and  $\Lambda$ -coalescents. This relation can be extended to a path-wise duality relation via a coupling of both processes that is based on the lookdown construction of [14], see section 2 in [8] and also [7] for details.

**Lemma 3.1** ([4];[13]). For fixed  $\phi \in \overline{B}(\mathcal{T}^p)$ ,  $p \ge 1$ , let  $H_{\phi}(\pi, \rho) \colon \mathscr{P}_{[p]} \times PM(\mathcal{T}) \to \mathbb{R}$  be given by

$$H_{\phi}(\pi,\rho) \coloneqq \langle \phi_{\pi}, \rho^{\otimes p} \rangle$$

where  $\phi_{\pi} \in \overline{B}(\mathcal{T}^{\#\pi})$  is constructed from  $\phi$  by identifying its input coordinates, say  $(a_1, \dots, a_p)$ , according to the blocks of  $\pi$ . Then we have

$$\mathbb{E}_{\rho_0}[\langle \phi_{\pi_0}, \rho_t^{\otimes p} \rangle] = \tilde{\mathbb{E}}_{\pi_0}[\langle \phi_{\Pi_t}, \rho_0^{\otimes p} \rangle]$$

whenever  $(\rho_t)_{t\geq 0}$  is a  $\Lambda$ -Fleming-Viot process under  $\mathbb{P}$  and  $(\Pi_t)_{t\geq 0}$  is a  $(p,\Lambda)$ -coalescent process under  $\tilde{\mathbb{P}}$ .

The case  $\Lambda = \frac{\sigma^2}{2} \delta_0$  corresponds to the standard Flemming-Viot process without mutation and of parameter  $\sigma$  which is dual to Kingman's coalescent and in which (9) becomes

$$\mathcal{R}F(\rho) = \frac{\sigma^2}{2} \int_{\mathcal{T}} \rho(da) (F''(\rho; a, a) - F(\rho)).$$

When  $\Lambda(0) = 0$ , the generator (9) can be written as, see [8],

$$\mathcal{R}F(\rho) = \int_{(0,1]} \frac{\Lambda(d\zeta)}{\zeta^2} \int_{\mathcal{T}} \rho(da) (F(\rho(1-\zeta) + \zeta\delta_a) - F(\rho)).$$

Combining both cases we obtain the following form of the generator

(10)  

$$\mathcal{R}F(\rho) = \frac{\sigma^2}{2} \int_{\mathcal{T}} \rho(da) (F''(\rho; a, a) - F(\rho)) + \int_{(0,1]} \frac{\Lambda(d\zeta)}{\zeta^2} \int_{\mathcal{T}} \rho(da) (F(\rho(1-\zeta) + \zeta\delta_a) - F(\rho)).$$

When  $\zeta^{-2}\Lambda(d\zeta)$  is finite, the above form of the generator gives the following picture for the dynamics of the process. It has jumps of the form  $\rho_{t-} \rightarrow \rho_{t-}(1-\zeta)+\zeta\delta_a$ at the atoms  $(t,\zeta)$  of a Poisson point process on  $\mathbb{R}_+ \times [0,1]$  with intensity  $dt \times \zeta^{-2}\Lambda(d\zeta)$ ; here the position a of the new atom of size  $\zeta$  is chosen randomly according to  $\rho_{t-}$ . After each jump, the process starts as an independent copy of a standard Flemming-Viot process of parameter  $\Lambda(\{0\})$  started at the new state  $\rho_t = \rho_{t-}(1-\zeta) + \zeta\delta_a$ .

An adaptation of the proof of Theorem 3 in [4] (to include the case of "pre-limit"  $\Lambda$ -measures not vanishing at {0}) yields the following weak-limit construction of the process for general finite measures  $\Lambda$ .

**Proposition 3.2.** For a sequence of measures  $\Lambda_n \xrightarrow[weakly]{n \to \infty} \Lambda$  such that  $\zeta^{-2}\Lambda_n(d\zeta)$  are finite measures in (0, 1], let  $(\rho_t^{(n)})_{t\geq 0}$  be corresponding  $\Lambda_n$ -Flemming-Viot processes started at  $\rho_0^{(n)}$ . If  $\rho_0^{(n)} \xrightarrow[d]{n \to \infty} \rho_0$  for the weak topology on  $\text{PM}(\mathcal{T})$ , then  $(\rho_t^{(n)})_{t\geq 0} \xrightarrow[d]{n \to \infty} (\rho_t)_{t\geq 0}$  for the Skorohod topology on  $D([0,\infty), \text{PM}(\mathcal{T}))$ , where  $(\rho_t)_{t\geq 0}$  is the  $\Lambda$ -Flemming-Viot process started at  $\rho_0$ .

### 3.3 The Dawson-Watanabe process and its Lamperti transformation

Here we introduce the Dawson-Watanabe process without mutation/spatial motion. This will suffice our applications further ahead; the interested reader can refer to [33] for a more general setting.

For fixed  $\sigma \in \mathbb{R}$ , the Dawson-Watanabe process without mutation can be defined as the unique continuous process  $(\mu_t)_{t\geq 0}$  on  $\mathbb{M}(\mathcal{T})$  such that for all  $\phi \in \overline{\mathbb{C}}(\mathcal{T})$  the process

$$M_t(\phi) = \langle \phi, \mu_t \rangle - \langle \phi, \mu_0 \rangle$$

is a martingale with quadratic variation

$$\left[M(\phi)\right]_t = \frac{\sigma^2}{2} \int_0^t \left<\phi^2, \mu_s\right> ds$$

Alternatively, it can be defined as the unique solution to the martingale problem for the operator

$$\mathcal{W}F(\mu) = \frac{\sigma^2}{2} \int_{\mathcal{T}} \mu(da) F^{\prime\prime}(\mu; a, a),$$

with domain  $\mathcal{D}(\mathcal{W})$  satisfying  $\{f(H(\mu)): H \in Pol(\mathbb{M}(\mathcal{T})), f \in C_0^{\infty}(\mathbb{R})\} \subset \mathcal{D}(\mathcal{W})$ . (see Corollary 2.23 in [17]).

Taking functions of the form  $H(\mu) = \langle 1, \mu \rangle$ , and  $f \in C_0^{\infty}(\mathbb{R})$  in this generator, it is easy to see that the total mass process  $(\|\mu_t\|)_{t\geq 0}$  has generator of the form  $\mathcal{K}f(x) = \frac{\sigma^2}{2}xf''(x)$  on the set  $\{f \in C_0^2(\mathbb{R}) : xf'' \in C([0, \infty])\} \subset \mathcal{D}(\mathcal{K})$  which conforms with the general form of the generator of a continuous positive 1-SS Markov process. By Theorem 5.1 in [30] the process  $(\|\mu_t\|)_{t\geq 0}$  is then a uniquely determined diffusion (Feller's diffusion), it is absorbed at 0 with probability 1 and, furthermore, by Theorem 4.1 therein the time-changed process  $(\log(\|\mu_{c_1(t)}\|))_{t\geq 0}$  is a continuous Lévy process with diffusion parameter  $\sigma$  and drift parameter  $-\sigma$ . The latter will be denoted d to be tracked as it will appear in the next sections.

On the other hand, it is well known from Theorem 1.1 i) in [8] (see also [32]) that the time-changed frequency process  $(\rho_t)_{t\geq 0}$  defined by  $\rho_t = \frac{\mu_{c_1(t)}}{\|\mu_{c_1(t)}\|}$  is a standard Fleming-Viot process of parameter  $\sigma$ .

We end this section with our first application of Theorem 2.2, which complements the latter observations, and which we will then generalize in Theorem 4.2.

**Proposition 3.3.** The process  $(\mu_t)_{t\geq 0}$  is a 1-SS Markov process. The time changed process  $(\nu_t)_{t\geq 0} = (\mu_{c_1(t)})_{t\geq 0}$  is SMH and its generator  $\mathcal{G}_{\sigma}^{(B)}$  has the form

(11) 
$$\mathcal{G}_{\sigma}^{(B)}F(\nu) := \|\nu\| \mathcal{W}F(\nu)$$

in the set

(12) 
$$D_1 := \{F \in \mathcal{D}(\mathcal{W}) \colon \|\cdot\| \, \mathcal{W}F(\cdot) \in \bar{\mathsf{B}}(\mathsf{M}(\mathcal{T}))\} \cup \{D \cap \mathcal{D}(\mathcal{W})\}.$$

Furthermore, the process  $(\rho_t, \xi_t)_{t\geq 0} = \left(\frac{\nu_t}{\|\nu_t\|}, \log(\|\nu_t\|)\right)_{t\geq 0}$  is a MAP with  $(\rho_t)_{t\geq 0}$  being a standard Fleming-Viot process of parameter  $\sigma$ , and  $(\xi_t)_{t\geq 0}$  a continuous Lévy process with diffusion parameter  $\sigma$  and drift parameter  $d = -\sigma$ .

*Proof.* We first check that  $(\nu_t)_{t\geq 0}$  is characterized by (11). Since  $(\|\mu_t\|)_{t\geq 0}$  is continuous and absorbed at 0, we get that

$$\inf_{s>0} \left\{ s \colon \|\mu_s\| = 0 \right\} = \inf_{a.s.\ s>0} \left\{ s \colon \int_0^s \frac{1}{\|\mu_s\|} ds = \infty \right\},$$

which implies that  $(\nu_t)_{t>0}$  is a solution to the martingale problem for  $\mathcal{G}_{\sigma}^{(B)}$  on the domain  $\{F \in \mathcal{D}(\mathcal{W}): \|\cdot\| \mathcal{W}F(\cdot) \in \bar{\mathbb{B}}(\mathbb{M}(\mathcal{T}))\}$  through a direct application of Theorem VI.1.3 in [16]. Thanks to the fact that the time-changed process  $(\|\nu_t\|)_{t\geq0}$  is the exponential of the continuous Lévy process given by  $\xi_t = dt + \sigma B_t$ , the domain can easily be extended to contain the set  $D \cap \mathcal{D}(\mathcal{W})$  by a mild adaptation of the proof of Theorem VI.1.3 in [16]. Indeed, we first note that  $N_t = F(\mu_t) - \int_0^t \mathcal{W}F(\mu_u)du$  is a martingale. Furthermore, for  $F \in D \cap \mathcal{D}(\mathcal{W})$ , we have the bound

$$\sup_{0 \le s \le t} |N_{c_1(s)}|^2 = \sup_{0 \le s \le t} \left| F(\mu_{c_1(s)}) - \int_0^{c_1(s)} WF(\mu_{c_1(u)}) du \right|^2$$
$$= \sup_{0 \le s \le t} \left| F(\nu_s) - \int_0^s \|\nu_u\| WF(\nu_u) du \right|^2 \le C(\sup_{0 \le s \le t} 1 + \dots + e^{p\xi_s})^2$$

for some  $C > 0, p \ge 1$ . By Doob's  $L^{2p}$  inequality applied to the submartingale  $(e^{2p\xi_t})_{t\ge 0}$ , the last line above is integrable. Using Hölder's inequality we also obtain

$$\lim_{T \to \infty} \mathbb{E}[|N_T|; c_1(t) > T] \le \mathbb{E}[\sup_{0 \le s \le t} |N_{c_1(s)}|^2]^{1/2} \mathbb{P}(c_1(t) > T)^{1/2} \to 0$$

where we have used  $c_1(t) < \infty$ . The optional sampling theorem (e.g. Theorem II.2.13 [16]) then implies that  $(N_{c_1(t)})_{t\geq 0}$  is a  $\mathscr{M}_{c_1(t)}$ -martingale.

On the other hand, the stated properties for  $(\rho_t, \xi_t)_{t\geq 0}$  will all follow from Theorem 2.2 once we prove that  $(\mu_t)_{t\geq 0}$  is 1-SS. The latter follows from Proposition 2.3 and the computation

$$\mathcal{WS}_a F(\mu) = \frac{\sigma^2}{2} \int_{\mathcal{T}} \mu(db) (\mathcal{S}_a F)^{\prime\prime}(\mu; b, b) = \frac{\sigma^2}{2} \int_{\mathcal{T}} \mu(db) a^2 F^{\prime\prime}(a\mu; b, b)$$

so that

$$S_{a^{-1}}\mathcal{W}S_aF(\mu) = a\frac{\sigma^2}{2}\int_{\mathcal{T}}\mu(db)F''(\mu;b,b) = a\mathcal{W}F(\mu).$$

### 4 The genealogy of self-similar populations

### 4.1 Main results and comments

We will now apply the generalized Lamperti transformation developed in section 2 for the conic subspace  $E = \mathbb{M}(\mathcal{T})$ . The first result is showing the existence of a Feller process with values in E enjoying the SMH property and having a generator of the form

(13) 
$$\mathcal{G}F(\nu) = \mathcal{G}_{\kappa}^{(D)}F(\nu) + \mathcal{G}_{\sigma}^{(B)}F(\nu) + \mathcal{G}_{\Lambda}^{(J)}F(\nu)$$

where  $\mathcal{G}_{\kappa}^{(D)}, \mathcal{G}_{\sigma}^{(B)}$  and  $\mathcal{G}_{\Lambda}^{(J)}$  are as in Theorem 1.1. The domain of the generator will be proved to contain the set

$$D_2 = \{ F \in D \cap D_1 \colon \mathcal{G}F \in \mathsf{C}(\mathsf{M}(\mathcal{T})) \},\$$

with D as in (8) and  $D_1$  as in (12).

**Theorem 4.1.** There exists a Feller process  $(\nu_t)_{t\geq 0}$  on  $\mathbb{M}(\mathcal{T})$  with generator of the form (13) in  $D_2$ . Furthermore, the process  $(\nu_t)_{t\geq 0}$  is SMH and is absorbed at 0.

The proof of the above theorem is split into several intermediary results. In section 4.2 we provide a Poissonian construction of the process  $(\nu_t)_{t\geq 0}$  when the measure  $\zeta^{-2}\Lambda(d\zeta)$  is finite; while in section 4.3 we extend the construction to any finite measure  $\Lambda$  through a weak limit and the convergence of the generators.

Having constructed the SMH process  $(\nu_t)_{t\geq 0}$ , our main result of this section is the following theorem which, on the one hand, generalizes Proposition 2 in [28] to  $\beta$ -stable measure-valued processes  $(\alpha > 1)$  and, on the other, re-frames Theorem 1 in [8] as a Lamperti transformation at least in the case  $\beta \in (1, 2]$  which comprises the intersection between branching and self-similar processes of index  $\alpha = \beta - 1$  (see Remark 2 below). Finally, the following theorem also describes the genealogy of populations whose total size evolves as a positive SS Markov process with non-negative jumps, through the well-known duality relationship between  $\Lambda$ -Fleming-Viot processes and  $\Lambda$ -coalescents [4] in the same spirit as [8] (see Lemma 3.1 and Remark 1). The core of the proof of the following Theorem 4.2 is already furnished by section 2.

**Theorem 4.2.** Let  $(\nu_t)_{t>0}$  be the SMH process constructed in Theorem 4.1.

- i) The process  $(\rho_t, \xi_t)_{t\geq 0}$  where  $\rho_t = \nu_t / \|\nu_t\|$  and  $\xi_t = \log(\|\nu_t\|)$  is a MAP. Furthermore,  $(\rho_t)_{t\geq 0}$  is a  $\Lambda$ -Fleming-Viot process, whereas  $(\xi_t)_{t\geq 0}$  is the Lévy process with triplet  $(d + \kappa, \sigma, \Pi)$  where  $\Pi(d\zeta)$  is the pushforward of the measure  $\zeta^{-2}\Lambda(d\zeta)$  under the transformation  $\zeta \to -\log(1-\zeta)$  on (0,1].
- ii) Let  $\alpha \in \mathbb{R}$  and recall the random time change  $\gamma_{\alpha}(t)$  of Theorem 2.2. The timechanged process  $(\mu_t)_{t>0} = (\nu_{\gamma_{\alpha}(t)})_{t>0}$  is the unique solution to

(14) 
$$\mu_t = \nu_{\int_0^t \|\mu_s\|^{-\alpha} ds}.$$

Furthermore, it is a  $\alpha$ -SS standard Markov process with generator of the form

(15) 
$$\mathcal{F}F(\mu) = \frac{1}{\|\mu\|^{\alpha}} \mathcal{G}F(\mu)$$

with domain containing  $\{F \in D_2 : \|\cdot\|^{-\alpha} \mathcal{G}F(\cdot) \in \overline{B}(\mathcal{M}(\mathcal{T}))\} \subset \mathcal{D}(\mathcal{F}).$ 

iii) Conversely, if  $(\mu_t)_{t\geq 0}$  is a standard Markov process with generator of the form (15). Then the process  $(\rho_t, \xi_t)_{t\geq 0}$  given by  $\rho_t = \mu_{c_{\alpha}(t)} / \|\mu_{c_{\alpha}(t)}\|$  and  $\xi_t = \log(\|\mu_{c_{\alpha}(t)}\|)$  is a MAP as in i). Furthermore,  $\nu_t = \rho_t e^{\xi_t}$  is the unique solution to

$$\nu_t = \mu_{\int_0^t \|\nu_s\|^\alpha ds}$$

and it is a standard Markov process with generator of the form (13).

Looking at  $\mathcal{G}_{\Lambda}^{(J)}$  in the generator of the process  $(\nu_t)_{t\geq 0}$  we observe that it has jumps of the form

$$\nu \to \nu + \|\nu\| x \delta_a$$

for some weight x > 0 and a location a for the new atom that is chosen according to the empirical distribution  $\nu / \|\nu\|$  before the jump. Since  $\|\nu\| x$ , the size of the new atom, depends on the total mass of the population, these processes will not be branching processes in general, nor will be their time-changed counterparts  $(\mu_t)_{t\geq 0}$ . However, the above form for the jumps of  $(\nu_t)_{t\geq 0}$  ensures that its corresponding frequency process is Markov with jumps

$$\rho = \frac{\nu}{\|\nu\|} \to \frac{\nu}{\|\nu\|} \left(\frac{1}{1+x}\right) + \frac{x}{1+x}\delta_a$$
$$= \rho\left(\frac{1}{1+x}\right) + \frac{x}{1+x}\delta_a.$$

The particular choice  $x = \zeta/(1-\zeta)$  ensures that the resulting dynamics are exactly those of (10) for FV processes. On the other hand, the dynamics of the time-changed process  $(\mu_t)_{t\geq 0}$  incorporate an additional dependency on the total mass by modulating the overall reproduction rate of the population according to  $\|\mu_t\|^{-\alpha}$  as seen in (15). By picking the right combination of the parameters  $\alpha$  and  $\Lambda \equiv \Lambda_{\alpha}$  one can show, via a simple computation on the generator (15), that these two sources of dependency on the total mass in fact cancel out. The resulting process  $(\mu_t)_{t\geq 0}$  is then a branching process, in particular a stable branching process (see Remark 2 below).

*Proof of Theorem 4.2.* Item i) will be proved along with the construction of the process  $(\nu_t)_{t>0}$  in sections 4.2 and 4.3.

Item ii) follows from first applying Theorem 2.2 to obtain that  $(\mu_t)_{t\geq 0}$  is a standard  $\alpha$ -SS Markov process. Now, observe that  $(\|\nu_t\|^{-\alpha})_{t\geq 0}$  is a.s. bounded on bounded time intervals since  $(\xi_t)_{t\geq 0}$  is a.s. bounded away from  $-\infty$  in bounded time intervals. The latter also implies that  $\inf\{t: \|\nu_t\|^{-\alpha} = 0\} = \infty = \gamma_{\alpha}(\infty)$ . Then by Theorem VI.1.1 in [16] the process  $(\mu_t)_{t\geq 0}$  is the unique solution to (14) and, furthermore, by Theorem VI.1.3 therein, it is a solution to the martingale problem for (15) on  $\{F \in D_2: \|\cdot\|^{-\alpha} \mathcal{G}F(\cdot) \in \bar{\mathbb{B}}(\mathcal{M}(\mathcal{T}))\}.$ 

Finally, iii) follows by first noting that  $(\|\mu_t\|)_{t\geq 0}$  is a positive  $\alpha$ -SS Markov process with non-negative jumps (see Remark 1). This implies that if  $\zeta_0^{\mu}$  is its absorption time at 0, then  $\mu_{\zeta_0^{\mu-}} \stackrel{=}{=} 0$ ; i.e.  $(\mu_t)_{t\geq 0}$  is absorbed continuously at 0. Then Lemmas 3.1 and 3.3 in [30] imply that  $c_{\alpha}(\infty) \stackrel{=}{=} \infty$ . In any case we have

$$\inf\{t: \|\mu_t\|^{\alpha} = 0\} = c_{\alpha}(\infty).$$

Then, a second application of Theorems VI.1.1 and VI.1.3 in [16] as before yields the result.  $\hfill \Box$ 

**Remark 1.** As in section 3.3, when taking functions of the form  $F(\mu) = f(\langle 1, \mu \rangle)$  in (15), we obtain that the total mass process  $(\|\mu_t\|)_{t\geq 0}$ , is a positive SS Markov process with generator

(16) 
$$\mathcal{K}f(x) = (d+\kappa)x^{1-\alpha}f'(x) + \frac{\sigma^2}{2}x^{2-\alpha}f''(x) + x^{-\alpha}\int_1^\infty [f(x\zeta) - f(x) - f'(x)\log(\zeta)\mathbb{1}_{|\log(\zeta)| < 1/2}]\Theta(d\zeta)$$

where  $\Theta(d\zeta)$  is the pushforward of the measure  $\zeta^{-2}\Lambda(d\zeta)$  under the transformation  $\zeta \to 1/(1-\zeta)$  (c.f. Theorem 6.1 in [30]).

**Remark 2.** Taking  $\sigma = 0$ , and  $\Lambda(d\zeta) = c\zeta^{1-\beta}(1-\zeta)^{\beta-1}d\zeta$  with c > 0 and  $\beta \in (1,2)$ ; and also

$$\kappa = -\int_0^1 \frac{\Lambda(d\zeta)}{\zeta^2} \left\{ \frac{\zeta}{1-\zeta} - \left( \left| \log(1-\zeta) \right| \mathbbm{1}_{\zeta < 1/2} \right) \right\},\,$$

in (15), we obtain

$$\mathcal{F}(\nu) = \frac{1}{\|\nu\|^{\beta-1}} \int_{\mathcal{T}} \frac{\nu(da)}{\|\nu\|} \int_{0}^{1} \frac{\Lambda(d\zeta)}{\zeta^{2}} \bigg\{ F\left(\nu + \|\nu\| \frac{\zeta}{1-\zeta} \delta_{a}\right) - F(\nu) - \|\nu\| \frac{\zeta}{1-\zeta} F'(\nu;a) \bigg\}.$$

The latter, after the change of variable  $h = \|\nu\| \frac{\zeta}{1-\zeta}$ , becomes

$$\mathcal{F}(\nu) = c \int_{\mathcal{T}} \nu(da) \int_0^\infty h^{-1-\beta} dh \bigg\{ F\left(\nu + h\delta_a\right) - F(\nu) - hF'(\nu;a) \bigg\},$$

the generator of a  $\beta$ -stable measure-valued branching process with  $\beta > 1$ . The case  $\beta = 2$  is covered by picking  $\Lambda(d\zeta) = c\delta_0(d\zeta)$ . Thus, Theorem 4.2 ii) generalizes Proposition 2 [28] to the measure-valued setting, while Theorem 4.2 iii) recovers Theorem 1 i) and ii) (for  $\beta > 1$ ) in [8].

**Remark 3.** Theorems 4.1 and 4.2 can be generalized in the two following directions using the same arguments in the proof, although details will not appear in this paper to avoid such burden in notation in our main results.

i) A general positive SS Markov process, see [3], can be attained for the total mass process (||μ<sub>t</sub>||)<sub>t≥0</sub> by adding negative jumps of the form x → xζ, 0 < ζ < 1, with intensity Θ<sup>-</sup>(dζ), the pushforward of a Lévy measure Π on (-∞,0) (i.e. a measure satisfying ∫<sup>0</sup><sub>-∞</sub>(ζ<sup>2</sup> ∧ 1)Π(dζ) < ∞) under the transformation ζ → e<sup>ζ</sup>. A simple way to do this is to allow for the underlying SMH process (ν<sub>t</sub>)<sub>t≥0</sub> to have jumps of the form ν → νζ with intensity Θ<sup>-</sup>(dζ). In this sense, a term of the form

$$\mathcal{G}_{\Theta^{-}}^{(J-)}F(\nu) = \int_{(0,1)} \Theta^{-}(d\zeta) \{F(\nu\zeta) - F(\nu)\}$$

is added to its generator. Adding negative jumps in this way does not change the dynamics of the corresponding frequency process, so that the arguments leading to the construction of the SMH process  $(\nu_t)_{t\geq 0}$  can be easily generalized in this direction.

ii) It is possible to extend our results to the  $\Xi$ -Fleming-Viot case with  $\int_{[0,1]^{\mathbb{N}}} \sum_{k=1}^{\infty} \zeta_k^2 \Xi(d\zeta) < \infty$ . In this case the generator of  $(\nu_t)_{t\geq 0}$  is given by updating the jumping part  $\mathcal{G}^{(J)}$  as

$$\begin{aligned} \mathcal{G}_{\Xi}^{(J)}F(\nu) &= \int_{(\mathcal{T})^{\mathbb{N}}} \left(\frac{\nu}{\|\nu\|}\right)^{\otimes \mathbb{N}} (da) \int_{(0,1]^{\mathbb{N}}} \Xi(d\zeta) \Big\{ F\left(\nu + \|\nu\| \sum_{i=1}^{\infty} \frac{\zeta_i}{1 - \|\zeta\|_{1_1}} \delta_{a_i}\right) - F(\nu) \\ &- \|\nu\| \left| \log(1 - \|\zeta\|_{1_1}) \right| \mathbbm{1}_{\|\zeta\|_{1_1} < 1/2} F'(\nu; a) \Big\}. \end{aligned}$$

Heuristically, jumps are of the form

$$\nu \to \nu + \sum_{k=1}^{\infty} \frac{\zeta_i}{1 - \|\zeta\|_{1_1}} \delta_{a_k}$$

where the positions of the new atoms  $(a_k)_{k\geq 0}$  are i.i.d with distribution  $\frac{\nu_{t-}}{\|\nu_{t-}\|}$ and the atom sizes  $\zeta = (\zeta_1, \zeta_2, \ldots)$ , satisfying  $\|\zeta\|_{1} < 1$ , arrive with intensity  $dt \otimes \Xi(d\zeta)$ .

The process  $(\rho_t)_{t\geq 0}$  becomes the  $\Xi$ -Flemming-Viot process and  $(\xi_t)_{t\geq 0}$  is a Lévy process with compensated jumps given by

$$\int_0^t \int_0^1 \left| \log(1 - \left\| \zeta \right\|_{1_1}) \right| \tilde{\mathcal{P}}(ds, d\zeta)$$

where  $\tilde{\mathcal{P}}(ds, d\zeta) = \mathcal{P}(ds, \zeta) - \mathbb{1}_{\|\zeta\|_{1_1} < 1/2} ds \otimes \Xi(d\zeta)$  and  $\mathcal{P}$  is a Poisson point process on  $\mathbb{R}^+ \times (0, 1)^{\mathbb{N}}$  with intensity  $ds \otimes \Xi(d\zeta)$ .

### **4.2** Poissonian construction for finite $\zeta^{-2}\Lambda(d\zeta)$

A simple way to think of the process  $(\nu_t)_{t\geq 0}$  when  $\zeta^{-2}\Lambda(d\zeta)$  is finite is as a timechanged and mass-scaled Dawson-Watanabe process to which atoms are added at times of an independent Poison point process (PPP). The size of such atoms will in general depend on the total population size at jump times.

Formally, for an initial condition  $\nu$ , let  $(\hat{\nu}_t)_{t\geq 0} = \left(\mu_{\gamma_1(t)}^{(DW)}\right)_{t\geq 0}$  where  $\left(\mu_t^{(DW)}\right)_{t\geq 0}$  is a standard  $\sigma$ -Dawson-Watanabe process started at  $\nu$ ; and  $\gamma_1(t) = \inf\left\{s \geq 0: \int_0^s \frac{1}{\|\mu_u^{(DW)}\|} du > t\right\}$ 

is the 1-SS Lamperti time change. We will work with several independent copies of  $(\hat{\nu}_t)_{t\geq 0}$  started at different initial measures  $\nu$ ; let us denote them by  $(\hat{\nu}_t(\nu))_{t\geq 0}$  when necessary.

Let

$$\hat{\kappa} \coloneqq \kappa - \int_{(0,1/2]} \left| \log(1-\zeta) \right| \frac{\Lambda(d\zeta)}{\zeta^2}$$

and let  $\mathcal{P}$  be a PPP with intensity  $dt \otimes \zeta^{-2} \Lambda(d\zeta)$  on  $\mathbb{R}_+ \times (0,1)$ . For t > 0 let  $(t_1, \zeta_1), \ldots, (t_K, \zeta_k)$  be the atoms of  $\mathcal{P}$  such that their first coordinate is less than t, ordered increasingly along the first (time) coordinate and set  $t_{K+1} = t$ . Then, conditionally on  $(t_i, \zeta_i)_{1 \leq i \leq K}$ , define  $\nu_s, s \in [0, t)$ , recursively as

(17) 
$$\nu_{s} \coloneqq \begin{cases} e^{\hat{\kappa}s}\hat{\nu}_{s}\left(\nu\right) & \text{if } 0 \le s < t_{1}, \\ e^{\hat{\kappa}(s-t_{i})}\hat{\nu}_{s-t_{i}}\left(\nu_{t_{i}-} + \|\nu_{t_{i}-}\|\frac{\zeta_{i}}{1-\zeta_{i}}\delta_{a_{i}}\right) & \text{if } t_{i} \le s < t_{i+1}, \end{cases}$$

where the processes  $(\hat{\nu}_t(\cdot))_{t\geq 0}$  that appear in each time interval  $[t_i, t_{i+1}]$  are all independent and the points  $(a_i)_{1\leq i\leq K}$  are chosen independently according to  $\nu_{t_i-}/ \|\nu_{t_i-}\|$ .

**Lemma 4.3.** The process  $(\nu_t)_{t\geq 0}$  is a standard SMH process with generator given by  $\mathcal{G}$  in (13) and domain  $\mathcal{D}(\mathcal{G}_{\sigma}^{(B)})$  containing  $D_1$ .

*Proof.* First, when  $\Lambda \equiv 0$ , i.e. for the Markov process  $(e^{\hat{\kappa}t}\hat{\nu}_t)_{t\geq 0}$ , a simple computation yields

$$\frac{d}{dt}\bigg|_{t=0} \mathbb{E}_{\nu}[F(e^{\hat{\kappa}t}\hat{\nu}_t)] = \mathcal{G}^{(D)}_{\hat{\kappa}}F(\nu) + \mathcal{G}^{(B)}_{\sigma}F(\nu).$$

To add the jumps, and compute the resulting generator, we use Theorem 2.4 in [36]. For this, we set (in their notation)  $\phi_t = C_{\Lambda} t$  with  $C_{\Lambda} \coloneqq \int_{[0,1]} \zeta^{-2} \Lambda(d\zeta)$  and the transition kernel  $K(\nu, \cdot)$  given by, for  $F \in \overline{B}(\mathbb{M}(\mathcal{T}))$ ,

$$K(\nu, F) = \begin{cases} \int_{(0,1]} \zeta^{-2} \frac{\Lambda(d\zeta)}{C_{\Lambda}} \int_{\mathcal{T}} \frac{\nu}{\|\nu\|} (da) F(\nu + \|\nu\| \frac{\zeta}{1-\zeta} \delta_a) & \text{whenever } \nu \neq 0, \\ F(0) & \text{otherwise.} \end{cases}$$

Then,  $\phi_t$  being of "Kac type", according to Theorem 2.4 (and Example 1) in [36], the generator of the process in (17) is given by

$$\begin{aligned} \frac{d}{dt} \bigg|_{t=0} \mathbb{E}[F(e^{\hat{\kappa}t}\hat{\nu}_t)] &= \mathcal{G}_{\hat{\kappa}}^{(D)}F(\nu) + \mathcal{G}_{\sigma}^{(B)}F(\nu) \\ &+ \int_{(0,1]} \zeta^{-2}\Lambda(d\zeta) \int_{\mathcal{T}} \frac{\nu}{\|\nu\|} (da) \left\{ F(\nu + \|\nu\| \frac{\zeta}{1-\zeta}\delta_a) - F(\nu) \right\} \\ &= \mathcal{G}_{\kappa}^{(D)}F(\nu) + \mathcal{G}_{\sigma}^{(B)}F(\nu) + \mathcal{G}_{\Lambda}^{(J)}F(\nu) \end{aligned}$$

with domain  $\mathcal{D}(\mathcal{G}_{\sigma}^{(B)})$ .

**Proposition 4.4.** The process  $(\rho_t, \xi_t)_{t\geq 0}$  where  $\rho_t = \nu_t / \|\nu_t\|$  and  $\xi_t = \log(\|\nu_t\|)$  is a MAP whose first coordinate is a  $(\frac{\sigma^2}{2}\delta_0 + \Lambda)$ -Fleming-Viot process, and its second coordinate is the Lévy process with triplet  $(d + \kappa, \sigma, \Pi)$  where  $\Pi(d\zeta)$  is the pushforward of the measure  $\zeta^{-2}\Lambda(d\zeta)$  under the transformation  $\zeta \to -\log(1-\zeta)$ . In particular  $\nu_t = \rho_t e^{\xi_t}$ , defines a Feller process.

We only provide a sketch of proof whose arguments can be formalized by further applications of Theorems 2.1 and 2.2 in [36]. We only provide the corresponding sample path analysis.

Sketch of proof. From the Poissonian construction of  $(\nu_t)_{t\geq 0}$ , and by direct calculations on its sample paths, we see that the normalized process defined by  $\rho_t = \nu_t / \|\nu_t\|$  has jumps of the form

$$\rho_{t-} = \frac{\nu_{t-}}{\|\nu_{t-}\|} \to \frac{\nu_{t-} + \|\nu_{t-}\| \left(\frac{\zeta}{1-\zeta}\right) \delta_a}{\|\nu_{t-}\| + \|\nu_{t-}\| \left(\frac{\zeta}{1-\zeta}\right)} = \frac{\nu_{t-} + \|\nu_{t-}\| \left(\frac{\zeta}{1-\zeta}\right) \delta_a}{\|\nu_{t-}\| \left(\frac{1}{1-\zeta}\right)} = \rho_{t-}(1-\zeta) + \zeta \delta_a$$

with intensity  $\zeta^{-2}\Lambda(d\zeta) \otimes \rho_{t-}(da)$ ; whereas, by Proposition 3.2, in-between jumps it evolves as a standard Fleming-Viot process of parameter  $\sigma$ .

Similarly,  $(\|\nu_t\|)_{t\geq 0}$  has jumps of the form

$$\|\nu_{t-}\| \to \|\nu_{t-}\| \left(1 + \frac{\zeta}{1-\zeta}\right),$$

so that  $\xi_t := \log(\|\nu_t\|)$  has jumps of the form

$$\xi_{t-} \to \xi_{t-} + \log\left(\frac{1}{1-\zeta}\right)$$

with intensity  $\zeta^{-2}\Lambda(d\zeta)$ . Also, in-between jumps, it has a drift  $\kappa$  and, by Proposition 3.2, an additional drift d and a Brownian component of parameter  $\sigma$ .

That  $(\rho_t, \xi_t)_{t\geq 0}$  is a MAP follows from  $(\nu_t)_{t\geq 0}$  being SMH (Lemma 4.3) and Proposition 2.1. Finally, the process  $(\rho_t, \xi_t)_{t\geq 0}$  is Feller since by Proposition 3.3 in [18] the  $\Lambda$ -Fleming-Viot process  $(\rho_t)_{t\geq 0}$  is Feller, as well as the Lévy process  $(\xi_t)_{t\geq 0}$ . The Feller property extends easily to the pair  $(\rho_t, \xi_t)_{t\geq 0}$ . Indeed, note that if  $\rho^{(k)} \to \rho$  implies  $\{\rho_t, \mathbb{P}_{\rho^{(k)}}\} \xrightarrow[d]{} \{\rho_t, \mathbb{P}_{\rho}\}$ , and also  $\xi^{(k)} \to \xi$  implies  $\{\xi_t, \mathbb{P}_{\xi^{(k)}}\} \xrightarrow[d]{} \{\xi_t, \mathbb{P}_{\xi^{(k)}}\}$ , then  $(\rho^{(k)}, \xi^{(k)}) \to (\rho, \xi)$  implies  $\{(\rho_t, \xi_t), \mathbb{P}_{(\rho^{(k)}, \xi^{(k)})}\} \xrightarrow[d]{} \{(\rho_t, \xi_t), \mathbb{P}_{(\rho^{(k)}, \xi^{(k)})}\}$ . Similarly, for the continuity in probability at t = 0, note that  $\rho_t \xrightarrow[e]{} \rho_0$  and  $\xi_t \xrightarrow[e]{} \phi_0$  imply  $(\rho_t, \xi_t) \xrightarrow[e]{} (\rho_0, \xi_0)$ . That  $(\nu_t)_{t\geq 0}$  is Feller now follows from the continuity of the mapping  $\phi^{-1}(\rho, \xi) = e^{\xi}\rho$  from  $\mathbb{PM}(\mathcal{T}) \times \mathbb{R}$  to  $\mathbb{M}(\mathcal{T})$ .

### 4.3 A Weak Limit and Martingale Problem

In this section we choose a particular sequence of processes  $\left\{ \left(\nu_t^{(n)}\right)_{t\geq 0} \right\}_n$  that will weakly approximate a process  $\left(\nu_t\right)_{t\geq 0}$  with generator  $\mathcal{G}$  as in (13), the latter will be proven to be a Feller process. Having proved that the limit process  $\left(\nu_t\right)_{t\geq 0}$  exists and is Feller, a general approximation theorem in terms of the convergence of the respective generators can be derived via Theorem 17.25 in [26]. In the following Proposition 4.5, and in order to simplify technical arguments, we only focus on the construction of the process  $\left(\nu_t\right)_{t\geq 0}$  for which we impose stronger conditions than just the convergence of the generators.

In the following we set  $\Lambda(\{0\}) = \frac{\sigma^2}{2}$ . Also, throughout this section, we will work with the jumping measure  $\mathbb{1}_{0 < \zeta \le 1/2} \Lambda(d\zeta)$  instead of  $\Lambda(d\zeta)$  for the limit process; while the approximating processes  $\left(\nu_t^{(n)}\right)_{t \ge 0}$  will have jumping measure

(18) 
$$\Lambda_n(d\zeta) = \mathbb{1}_{1/n < \zeta < 1/2} \Lambda(d\zeta).$$

Removing the atoms in which  $\zeta > 1/2$  ensures that the jumps of  $\left( \left\| \nu_t^{(n)} \right\| \right)_{t \ge 0}$  are bounded, which in turn simplifies the proof of tightness for the family  $\left\{ \left( \nu_t^{(n)} \right)_{t \ge 0} \right\}_n$ . This imposes no loss of generality for the construction of the process  $(\nu_t)_{t \ge 0}$  with generator of the general form  $\mathcal{G}$  since the finitely-many large jumps that occur with intensity  $\mathbbm{1}_{\zeta>1/2}\Lambda(d\zeta)$  can always be "added back" via the same argument as in Lemma 4.3; whilst ensuring that the resulting frequency process  $(\rho_t)_{t \ge 0}$  is the corresponding  $\Lambda$ -Felming-Viot process and the logarithm of the norm  $(\xi_t)_{t \ge 0}$  is the corresponding Lévy process as in Proposition 4.4.

**Proposition 4.5.** Let  $\left(\nu_t^{(n)}\right)_{t\geq 0}$  be constructed as in section 4.2 with  $\Lambda_n$  as in (18) where, furthermore,  $\nu_0^{(n)} \xrightarrow[d]{\to} \nu_0$  and  $\sup_n \mathbb{E}[\left\|\nu_0^{(n)}\right\|^p] < \infty$  for every  $p \geq 1$ . Let  $\left(\rho_t^{(n)}, \xi_t^{(n)}\right)_{t\geq 0}$  be the corresponding MAPs. Then, as  $n \to \infty$ ,

$$\left(\rho_t^{(n)}, \xi_t^{(n)}, \nu_t^{(n)}\right)_{t \ge 0} \Rightarrow \left(\rho_t, \xi_t, \nu_t\right)_{t \ge 0}$$

in the Skorohod topology on  $D([0,\infty); \mathsf{PM}(\mathcal{T}) \times \mathbb{R} \times \mathsf{M}(\mathcal{T}))$ , where

$$(\nu_t)_{t\geq 0} = \left(e^{\xi_t}\rho_t\right)_{t\geq 0}$$

and  $(\rho_t)_{t\geq 0}$  is a  $\Lambda$ -Fleming-Viot process, and  $(\xi_t)_{t\geq 0}$  a Lévy process with triplet  $(d + \kappa, \sigma, \Pi)$  where  $\Pi(d\zeta)$  is the pushforward of the measure  $\zeta^{-2}\Lambda(d\zeta)$  under the transformation  $\zeta \to -\log(1-\zeta)$ . Furthermore, the process  $(\nu_t)_{t\geq 0}$  is Feller with generator of the form  $\mathcal{G}$  in  $D_2$ .

*Proof.* We proceed in multiple steps: proving existence of solutions to the  $\mathcal{G}$ -martingale problem through tightness of the family of processes  $\left\{ \left( \nu_t^{(n)} \right)_{t \geq 0} \right\}_n$  in the Skorohod topology for  $D([0,\infty); \mathbb{M}(\mathcal{T}))$ , and the convergence of the approximating generators  $\mathcal{G}_n$  to  $\mathcal{G}$ ; and then identifying the limit and showing that it is Feller. The proof of tightness rests on the well-known Aldous-Rebolledo criterion (e.g. Theorem 1.17 in [15]), whereas the properties of the limiting process follow through an analysis of  $(\rho_t)_{t\geq 0}$  and  $(\xi_t)_{t\geq 0}$  separately.

Step 1: Tightness. The proof rests on Theorem III.9.1 in [16]. Let us first prove that

(19) 
$$\lim_{r \to \infty} \limsup_{n \ge 1} \mathbb{P}\left(\sup_{0 \le t \le T} \left\|\nu_t^{(n)}\right\| \ge r\right) = 0$$

which implies the compact containment condition (eq. (9.1) therein) for our processes since,  $\mathcal{T}$  being compact, the space  $\mathbb{M}(\mathcal{T})$  is locally compact. In fact the balls of radius  $r, \mathbb{M}_r(\mathcal{T}) = \{\mu \in \mathbb{M}(\mathcal{T}) : \langle 1, \mu \rangle \leq r\}$ , are compact (see e.g. Theorem 1.14 in [31]).

$$\begin{split} r, & \mathbb{M}_r\left(\mathcal{T}\right) = \{\mu \in \mathbb{M}\left(\mathcal{T}\right): \ \langle 1, \mu \rangle \leq r\}, \text{ are compact (see e.g. Theorem 1.14 in [31])}.\\ & \text{By Proposition 4.4 we have } \left\|\nu_t^{(n)}\right\| = e^{\xi_t^{(n)}} \text{ where } \left(\xi_t^{(n)}\right)_{t\geq 0} \text{ is the Lévy process } \end{split}$$

$$\xi_t^{(n)} = \log(\left\|\nu_0^{(n)}\right\|) + (\kappa + d)t + M_t^{(n)}.$$

Here  $\left(M_t^{(n)}\right)_{t\geq 0}$  is its martingale component, given by

$$M_t^{(n)} = \sigma B_t^{(n)} + \int_0^t \int_{1/n < \zeta < 1/2} |\log(1-\zeta)| \,\tilde{\mathcal{P}}^{(n)}(ds, d\zeta),$$

where  $(B_t^{(n)})_{t\geq 0}$  is a brownian motion and  $\tilde{\mathcal{P}}^{(n)}$  is a compensated Poisson random measure on  $\mathbb{R}_+ \times [1/n, 1/2]$  of intensity  $ds \times \zeta^{-2} \Lambda(d\zeta)$ . Markov's inequality followed by Hölder's inequality yield, for any p, q with 1/p + 1/q = 1,

(20) 
$$\mathbb{P}\left(\sup_{0\leq t\leq T} \left\|\nu_t^{(n)}\right\| \geq r\right) \leq \frac{e^{|\kappa+d|T}}{r} \mathbb{E}\left[\left\|\nu_0^{(n)}\right\| \sup_{0\leq t\leq T} e^{M_t^{(n)}}\right] \leq \frac{e^{|\kappa+d|T}}{r} \mathbb{E}\left[\left\|\nu_0^{(n)}\right\|^q\right]^{1/q} \mathbb{E}\left[\sup_{0\leq t\leq T} e^{pM_t^{(n)}}\right]^{1/p}.$$

The  $L^q$ -norm of  $\|\nu_0^{(n)}\|$  above is uniformly bounded on n by hypothesis. We now bound, also uniformly on n, the second expectation in the r.h.s. above. Then taking  $r \to \infty$ in the last line above will yield (19). By Doob's maximal inequality applied to the submartingale  $\left(e^{pM_t^{(n)}}\right)_{t\geq 0}$  we have for some constant  $C_p$ , and plugging Campbell's formula in the second inequality below,

(21) 
$$\mathbb{E}[\sup_{0 \le t \le T} e^{pM_t^{(n)}}] \le C_p \mathbb{E}[e^{pM_T^{(n)}}]$$
$$\le C_p e^{Tp^2 \frac{\sigma^2}{2}} \exp\left\{T \int_{1/n}^{1/2} \frac{\Lambda(d\zeta)}{\zeta^2} \left\{e^{-p\log(1-\zeta)} - 1 + p\log(1-\zeta)\right\}\right\}.$$

Since the integrand above is of order  $\mathcal{O}(\zeta^2)$  as  $\zeta \to 0$ , we have

$$\int_0^{1/2} \frac{\Lambda(d\zeta)}{\zeta^2} \left\{ e^{-p\log(1-\zeta)} - 1 + p\log(1-\zeta) \right\} < \infty,$$

which gives a bound that is uniform on n for (21).

In view of Theorem III.9.1 in [16], and the fact that the polynomials  $Pol(M(\mathcal{T}))$  are dense in  $\overline{C}(M(\mathcal{T}))$  for the topology of uniform convergence in compact sets; it remains to prove that  $\left\{ \left( F(\nu_t^{(n)}) \right)_{t \ge 0} \right\}_n$  is relatively compact for every  $F \in \operatorname{Pol}(\mathbb{M}(\mathcal{T})) \subset D$ , where we recall that D is as in (8). For the latter, in turn, we will prove the conditions of the Aldous-Rebolledo criterion.

First, the tightness of  $\{F(\nu_t^{(n)})\}_n$  for fixed t follows directly from (19). Second, by Lemma 4.3,  $N_t^{(n)} \coloneqq F(\nu_t^{(n)}) - \int_0^t \mathcal{G}_n F(\nu_s^{(n)}) ds$  defines a martingale so that the finite variation process of  $\left(F(\nu_t^{(n)})\right)_{t\geq 0}$  is given by

$$V_t^{(n)} = \int_0^t \mathcal{G}_n F(\nu_s^{(n)}) ds.$$

On the other hand, the compensator of the process  $\left((N_t^{(n)})^2\right)_{t>0}$  is given by

$$\left[N^{(n)}\right]_t = \int_0^t \mathbf{\Gamma}_n F(\nu_s^{(n)}) ds$$

where  $\Gamma_n$  is the carré du champ operator associated to  $\mathcal{G}_n$ . The latter is given by

$$\Gamma_n F = \mathcal{G}_n F^2 - 2F \mathcal{G}_n F$$

(see e.g. section 1.2.2 in [24]). The remaining conditions of the Aldous-Rebolledo criterion on  $(V_t^{(n)})_{t\geq 0}$  and  $([N^{(n)}]_t)_{t\geq 0}$  (which are given by (23) and (24) below) will follow once we prove that for any  $F \in \mathsf{Pol}(\mathsf{M}(\mathcal{T})), \delta > 0$ , and any stopping time  $0 \le \tau_n \le T,$ 

(22) 
$$\limsup_{n \to \infty} \sup_{0 \le \theta \le \delta} \mathbb{E} \left[ \int_{\tau_n}^{\tau_n + \theta} \left( \left| F(\nu_s^{(n)}) \right| \lor 1 \right) \left| \mathcal{G}_n F(\nu_s^{(n)}) \right| ds \right] \le \delta C$$

for some C > 0 depending only on F. Indeed, by Markov's inequality we obtain, on the one hand,

$$\begin{split} &\limsup_{n \to \infty} \sup_{0 \le \theta \le \delta} \mathbb{P} \left( \left| V_{\tau_n + \theta}^{(n)} - V_{\tau_n}^{(n)} \right| > \epsilon \right) \\ &\le \limsup_{n \to \infty} \sup_{0 \le \theta \le \delta} \mathbb{E} \left[ \int_{\tau_n}^{\tau_n + \theta} \left| \mathcal{G}_n F(\nu_s^{(n)}) \right| ds \right] \le \frac{\delta C}{\epsilon}. \end{split}$$

So that, taking  $\delta = \epsilon^2 / C$ , (22) implies

(23) 
$$\sup_{n \ge n_0} \sup_{0 \le \theta \le \delta} \mathbb{P}\left[ \left| V_{\tau_n + \theta}^{(n)} - V_{\tau_n}^{(n)} \right| > \epsilon \right] \le \epsilon$$

On the other hand, for any  $\delta > 0$ ,

$$\mathbb{E}\left[\left|\left[N^{(n)}\right]_{\tau_{n}+\delta}-\left[N^{(n)}\right]_{\tau_{n}}\right|\right]$$
  
$$\leq \mathbb{E}\left[\int_{\tau_{n}}^{\tau_{n}+\delta}\left|\mathcal{G}_{n}F^{2}(\nu_{s}^{(n)})\right|ds\right]+2\mathbb{E}\left[\int_{\tau_{n}}^{\tau_{n}+\delta}\left|F(\nu_{s}^{(n)})\mathcal{G}_{n}F(\nu_{s}^{(n)})\right|ds\right],$$

so that now (22) applied to each term in the r.h.s. above, and choosing  $\delta$  adequately, yields

(24) 
$$\sup_{n \ge n_0} \sup_{0 \le \theta \le \delta} \mathbb{P}\left[ \left| \left[ N^{(n)} \right]_{\tau_n + \theta} - \left[ N^{(n)} \right]_{\tau_n} \right| > \epsilon \right] \le \epsilon$$

again through Markov's inequality.

Let us then prove (22). On the one hand, by (8), the diffusion and drift terms of  $\mathcal{G}_n F$  are uniformly bounded on  $n \geq 1$  by

$$\begin{aligned} \left| \mathcal{G}_{\sigma}^{(B)} F(\nu) \right| + \left| \mathcal{G}_{\kappa}^{(D)} F(\nu) \right| &\leq \|\nu\| \int_{\mathcal{T}} \nu(da) \frac{\sigma^2}{2} \left| F''(\nu; a, a) \right| + \int_{\mathcal{T}} \nu(da) \kappa \left| F'(\nu; a) \right| \\ (25) &\leq C_1 \left( 1 + \dots + \|\nu\|^{p_1} \right) \end{aligned}$$

for some  $C_1 > 0$  and  $p_1 \ge 1$ . On the other hand, for the jump component  $\mathcal{G}_{\Lambda_n}^{(J)} F$  built from the measure  $\Lambda_n$  defined in (18), we have

(26) 
$$\left\|\Lambda_n(d\zeta) - \mathbb{1}_{0 < \zeta \le 1/2} \Lambda(d\zeta)\right\|_{TV} = \Lambda([0, \frac{1}{n})) \xrightarrow{n \to \infty} 0.$$

Fix a measure  $\nu$  and regard  $F(\nu + \|\nu\| \frac{\zeta}{1-\zeta} \delta_a)$  as a function of  $\zeta$ . Then writing  $\psi(\zeta) = \frac{\zeta}{1-\zeta}$  so that  $\psi'(\zeta) = \frac{1}{(1-\zeta)^2}$  and  $\psi''(\zeta) = \frac{2}{(1-\zeta)^3}$ , we have

$$\frac{d}{d\zeta}F(\nu + \|\nu\|\psi(\zeta)\delta_a) 
= \lim_{h \to 0} \frac{F(\nu + \|\nu\|\psi(\zeta)\delta_a + \|\nu\|(\psi(\zeta + h) - \psi(\zeta))\delta_a) - F(\nu + \|\nu\|\psi(\zeta)\delta_a)}{\|\nu\|(\psi(\zeta + h) - \psi(\zeta))} 
\times \lim_{h \to 0} \|\nu\|\frac{\psi(\zeta + h) - \psi(\zeta)}{h} 
= \|\nu\|F'(\nu + \|\nu\|\psi(\zeta)\delta_a; a)\psi'(\zeta),$$

and, similarly,

$$\frac{d}{d\zeta^2} F(\nu + \|\nu\| \psi(\zeta)\delta_a) = \|\nu\|^2 F''(\nu + \psi(\zeta)\delta_a; a, a)(\psi'(\zeta))^2 + \|\nu\| F'(\nu + \psi(\zeta)\delta_a; a)\psi''(\zeta).$$

Taylor's expansion for  $\zeta$  near 0 and the assumption  $F \in D$  give, for some constant  $C_2 > 0$  and some  $p_2 \ge 1$ ,

$$\left| F\left(\nu + \|\nu\| \frac{\zeta}{1-\zeta} \delta_a \right) - F(\nu) - \|\nu\| \zeta F'(\nu; a) \right| \le C_2 \left(1 + \dots + \|\nu\|^{p_2}\right) \frac{\zeta^2}{(1-\zeta)^4}.$$

Furthermore, using that

$$\forall 0 < \zeta < 1, \quad |\zeta - |\log(1 - \zeta)|| = \sum_{k=2}^{\infty} \frac{\zeta^k}{k} \le \frac{\zeta^2}{1 - \zeta}$$

which is continuous and bounded on  $\zeta \in [0, 1/2)$ , we see that the function of  $\zeta$ 

$$\mathbb{1}_{\zeta\in(0,1]}\left\langle\frac{F\left(\nu+\|\nu\|\frac{\zeta}{1-\zeta}\delta_a\right)-F(\nu)-\|\nu\|\left|\log(1-\zeta)\right|\mathbb{1}_{\zeta<1/2}F'(\nu;a)}{\zeta^2},\frac{\nu(da)}{\|\nu\|}\right\rangle$$

is bounded uniformly on [0, 1/2] by  $C_3(1 + \cdots + \|\nu\|^{p_2})$  for some  $C_3 > 0$  depending only on F. Thus (26) yields

(27) 
$$\left|\mathcal{G}_{\Lambda_n}^{(J)}F(\nu) - \mathcal{G}_{\Lambda}^{(J)}F(\nu)\right| \le C_3 \left(1 + \dots + \|\nu\|^{p_2}\right) \|\Lambda_n(d\zeta) - \Lambda(d\zeta)\|_{TV}$$

and, in particular,

(28) 
$$\left|\mathcal{G}_{\Lambda_n}^{(J)}F(\nu)\right| \leq C_4 \left(1 + \dots + \|\nu\|^{p_2}\right)$$

for some  $C_4$ . Putting together (25) and (28), and also using that  $F(\nu)$  satisfies the same type of inequality (by (8)), we obtain for some C > 0 and  $p \ge 1$ , and for any stopping time  $0 \le \tau_n \le T$  and  $0 \le \theta \le \delta < 1$ ,

(29) 
$$\mathbb{E}\left[\int_{\tau_n}^{\tau_n+\theta} \left(\left|F(\nu_s^{(n)})\right| \vee 1\right) \left|\mathcal{G}_n F(\nu_s^{(n)})\right| ds\right] \le C\delta \mathbb{E}\left[\sup_{t \in [0,T+\delta]} 1 + \dots + e^{p\xi_t^{(n)}}\right]$$

where we have again used  $\left(\left\|\nu_t^{(n)}\right\|\right)_{t\geq 0} = \left(e^{\xi_t^{(n)}}\right)_{t\geq 0}$  from Proposition 4.4. It remains to prove that the expectation in the r.h.s. above is uniformly bounded on  $n\geq 1$ . But this follows from (20) and (21).

Step 2: Limiting martingale problem. Assume that, along a subsequence  $\{n_i\}_{i\geq 1}$  (that we will denote by *n* to ease notations) we have

$$\left(\nu_t^{(n)}\right)_{t\geq 0} \stackrel{n\to\infty}{\xrightarrow[d]{\to}} \left(\nu_t\right)_{t\geq 0}$$

in  $D([0,\infty), \mathbb{M}(\mathcal{T}))$  for some process  $(\nu_t)_{t\geq 0}$ . We now prove that  $(\nu_t)_{t\geq 0}$  must solve the martingale problem for (13) in the domain  $\{F \in D : \mathcal{G}F \in C(\mathbb{M}(\mathcal{T}))\}$ .

We prove the latter via a mild adaptation of the proof of Lemma IV.5.1 in [16] to the case where F and  $\mathcal{G}F$  may not be bounded (but still continuous), and also where

the convergence of the generators  $\mathcal{G}_n$  to  $\mathcal{G}$  is uniform but only on compact sets (taking advantage of the fact that we have already proved the compact containment condition (19)). We only prove that the key equation (IV.5.3) therein (eq. (32) below) also holds in our setting.

Let  $h_i, 1 \leq i \leq k$ , be functions in  $\overline{C}(M(\mathcal{T}))$ . Also let  $0 \leq s_i \leq s < t$  all belong to the set  $\{u: \nu_u = \nu_{u-} \text{ a.s.}\}$ . Then, for  $F \in D$  such that  $\mathcal{G}F \in C(M(\mathcal{T}))$ , we have, by the continuous mapping theorem,

$$(F(\nu_t^{(n)}) - F(\nu_s^{(n)}) - \int_s^t \mathcal{G}F(\nu_u^{(n)}) du) \prod_i h_i(\nu_{s_i}^{(n)})$$
  
$$\stackrel{n \to \infty}{\longrightarrow} (F(\nu_t) - F(\nu_s) - \int_s^t \mathcal{G}F(\nu_u) du) \prod_i h_i(\nu_{s_i}).$$

The corresponding convergence in expectation follows from Theorem 3.5 in [6] and the fact that the random variables in the first line above are uniformly integrable. To verify the uniform integrability, observe that, since  $F \in D$ , the first line above is bounded by

(30) 
$$\left| (F(\nu_t^{(n)}) - F(\nu_s^{(n)}) - \int_s^t \mathcal{G}F(\nu_u^{(n)}) du) \prod_{i=1}^k h_i(\nu_{s_i}^{(n)}) \right|$$
$$\leq C(\sup_{s \le u \le t} 1 + \dots + \left\| \nu_u^{(n)} \right\|^p) \prod_{i=1}^k \|h_i\|_{\infty}$$

for some  $C, p \ge 0$  (where we have used also (28)). That the r.h.s. is uniformly integrable, in turn, is a simple consequence of Theorem 3.6 in [6] together with the convergence

$$\sup_{s \le u \le t} \left\| \nu_u^{(n)} \right\|^p \stackrel{n \to \infty}{\xrightarrow[]{d}{\longrightarrow}} \sup_{s \le u \le t} \left\| \nu_u \right\|^p,$$

and the bound in (21). Indeed, the corresponding a.s. convergence ensured by Skorohod representation theorem, and the bound in (21), together yield the required convergence in expectation

$$\mathbb{E}\left[\sup_{s\leq u\leq t}\left\|\nu_{u}^{(n)}\right\|^{p}\right]\overset{n\to\infty}{\longrightarrow}\mathbb{E}\left[\sup_{s\leq u\leq t}\left\|\nu_{u}\right\|^{p}\right].$$

We thus conclude

(31) 
$$\mathbb{E}[(F(\nu_t) - F(\nu_s) - \int_s^t \mathcal{G}F(\nu_u) du) \prod_{i=1}^k h_i(\nu_{s_i})] = \lim_{n \to \infty} \mathbb{E}[(F(\nu_t^{(n)}) - F(\nu_s^{(n)}) - \int_s^t \mathcal{G}F(\nu_u^{(n)}) du) \prod_{i=1}^k h_i(\nu_{s_i}^{(n)})].$$

We now approximate  $\mathcal{G}$  by  $\mathcal{G}_n$  in the above limit. Observe that, by (27), and since the other two components of  $\mathcal{G}_n$  are constant on n, we have

$$|\mathcal{G}_n F(\nu) - \mathcal{G}F(\nu)| \le C_3 \left(1 + \dots + \|\nu\|^{p_2}\right) \Lambda((0, 1/n)) \xrightarrow{n \to \infty} 0$$

uniformly for  $\nu$  on the compact sets  $M_r$ ,  $r \ge 0$ . Plugging the above bound into (31), together with the bound in (30) and the uniform integrability of  $\sup_{s\le u\le t} \left\| \nu_u^{(n)} \right\|^p$ ,

easily yield

(32) 
$$\mathbb{E}[(F(\nu_t) - F(\nu_s) - \int_s^t \mathcal{G}F(\nu_u) du) \prod_{i=1}^k h_i(\nu_{s_i})] = \lim_{n \to \infty} \mathbb{E}[(F(\nu_t^{(n)}) - F(\nu_s^{(n)}) - \int_s^t \mathcal{G}_n F(\nu_u^{(n)}) du) \prod_{i=1}^k h_i(\nu_{s_i}^{(n)})] = 0,$$

where we have used Lemma 4.3 in the last equality.

Step 3: Identification of the limit and Feller property. We first show that the sequence  $\left\{\left(\log\left(\left\|\nu_{t}^{(n)}\right\|\right)\right)_{t\geq0}\right\}_{n}$  converges in distribution towards the Lévy process  $(\xi_{t})_{t\geq0}$ , and that the sequence  $\left\{\left(\rho_{t}^{(n)}\right)_{t\geq0}\right\}_{n}$  converges in distribution towards the  $\Lambda$ -Fleming-Viot process  $(\rho_{t})_{t\geq0}$ . Indeed, this follows by the tightness of the family  $\left\{\left(\nu_{t}^{(n)}\right)_{t\geq0}\right\}_{n}$  and the continuous mapping theorem on the one hand, and on the other by identifying the limiting processes along subsequences through martingale problem characterizations. The latter is done by taking functions of the form  $F(\nu) = f(\log(\|\nu\|))$  with  $f \in C_{0}^{2}(\mathbb{R})$ , and  $F_{\phi}(\nu) = \left\langle \phi, \left(\frac{\nu}{\|\nu\|}\right)^{\otimes p} \right\rangle$  with  $\phi \in \overline{\mathbb{B}}(\mathcal{T}^{p})$ , respectively in (13) and using the uniqueness of solutions to the respective martingale problems of the Lévy and the  $\Lambda$ -Fleming-Viot processes (one could also use classical Lévy processes criteria for weak convergence such as Theorem VII.3.4 in [23] on the one hand, and on the other Proposition 3.2 for the Fleming-Viot part). Indeed, we check the computations only for the generator of the frequency process, which will coincide with that of the  $\Lambda$ -Fleming-Viot process. For

$$F_{\phi}(\nu) = G_{\phi}\left(\frac{\nu}{\|\nu\|}\right), \quad G_{\phi}(\rho) = \left\langle \phi, \rho^{\otimes p} \right\rangle,$$

we have, on the one hand,

$$F_{\phi}'(\nu;a) = \frac{1}{\|\nu\|^{2p}} \left( \sum_{i=1}^{p} \left\{ \left\langle \phi, \nu^{\otimes i} \delta_{a} \nu^{\otimes p-i-1} \right\rangle \|\nu\|^{p} \right\} - p \|\nu\|^{p-1} \left\langle \phi, \nu^{\otimes p} \right\rangle \right)$$
$$= \frac{1}{\|\nu\|} \sum_{i=1}^{p} \left\{ \frac{\left\langle \phi, \nu^{\otimes i} \delta_{a} \nu^{\otimes p-i-1} \right\rangle}{\|\nu\|^{p-1}} - \frac{\left\langle \phi, \nu^{\otimes p} \right\rangle}{\|\nu\|^{p}} \right\}$$
$$(33) \qquad = \frac{1}{\|\nu\|} \sum_{i=1}^{p} \left\{ F_{\phi_{i}^{(a)}}(\nu) - F_{\phi}(\nu) \right\}$$

where we have written

$$F_{\phi_i^{(a)}}(\nu) = \frac{\left\langle \phi_i^{(a)}, \nu^{\otimes p} \right\rangle}{\|\nu\|^p} = \frac{\left\langle \phi, \nu^{\otimes i} \delta_a \nu^{\otimes p-i-1} \right\rangle}{\|\nu\|^{p-1}}$$

with  $\phi_i^{(a)}(a_1, \dots, a_p) = \phi(a_1, \dots, a_{i-1}, a, a_{i+1}, \dots, a_p)$ . More generally, for a subset  $J \subset [p]$  the function  $\phi_J^{(a)} \in \overline{\mathbb{B}}(\mathcal{T}^p)$  will denote the function constructed from  $\phi$  by forcing all the coordinates in J to have the value  $a \in \mathcal{T}$ , and evaluating  $\phi$  at the new location.

From the above we obtain

(34) 
$$\int_{\mathcal{T}} \frac{\nu(da)}{\|\nu\|} \|\nu\| F_{\phi}'(\nu; a) = \sum_{i=1}^{p} \left\{ \frac{\langle \phi, \nu^{\otimes p} \rangle}{\|\nu\|^{p}} - \frac{\langle \phi, \nu^{\otimes p} \rangle}{\|\nu\|^{p}} \right\} = 0.$$

On the other hand, writing  $\mathcal{P}(a, \ell, \nu, p)$  for the set of all permutations of  $\ell$  copies of the measure  $\delta_a$  together with  $p - \ell$  copies of  $\nu$ , we have

$$F_{\phi}(\nu + \|\nu\| \frac{\zeta}{1-\zeta} \delta_{a}) - F_{\phi}(\nu)$$

$$= \frac{\langle \phi, \nu^{\otimes p} \rangle}{\|\nu\|^{p}} \left( (1-\zeta)^{p} - 1 \right) + \sum_{\ell=1}^{p} \sum_{\sigma \in \mathcal{P}(a,\ell,\nu,p)} \frac{\langle \phi, \otimes_{i=1}^{p} \sigma_{i} \rangle}{\|\nu\|^{p} (1-\zeta)^{-p}} \left( \frac{\|\nu\| \zeta}{1-\zeta} \right)^{\ell}$$

$$= \frac{\langle \phi, \nu^{\otimes p} \rangle}{\|\nu\|^{p}} \left( (1-\zeta)^{p} - 1 \right) + \sum_{\ell=1}^{p} \sum_{\sigma \in \mathcal{P}(a,\ell,\nu,p)} \frac{\langle \phi, \otimes_{i=1}^{p} \sigma_{i} \rangle}{\|\nu\|^{p-\ell}} (1-\zeta)^{p-\ell} \zeta^{\ell}$$

$$= \sum_{\ell=0}^{p} (1-\zeta)^{p-\ell} \zeta^{\ell} \sum_{\sigma \in \mathcal{P}(a,\ell,\nu,p)} \left\{ \frac{\langle \phi, \otimes_{i=1}^{p} \sigma_{i} \rangle}{\|\nu\|^{p-\ell}} - \frac{\langle \phi, \nu^{\otimes p} \rangle}{\|\nu\|^{p}} \right\}.$$

Observe that for  $\ell = 0$  we have

$$\int_{\mathcal{T}} \frac{\nu(da)}{\|\nu\|} \frac{\left\langle \phi, \nu^{\otimes p} \right\rangle}{\|\nu\|^{p}} - \frac{\left\langle \phi, \nu^{\otimes p} \right\rangle}{\|\nu\|^{p}} = 0,$$

whereas, by (33) and (34), the  $\ell = 1$  term similarly vanishes in  $\mathcal{G}_{\Lambda}^{(J)}$ . Then, substituting above, we obtain

$$\int_{\mathcal{T}} \frac{\nu(da)}{\|\nu\|} \left\{ F_{\phi}(\nu + \|\nu\| \frac{\zeta}{1-\zeta} \delta_{a}) - F_{\phi}(\nu) \right\}$$
$$= \sum_{\ell=2}^{p} (1-\zeta)^{p-\ell} \zeta^{\ell} \sum_{\sigma \in \mathcal{P}(a,\ell,\nu,p)} \left\{ \frac{\langle \phi, \otimes_{i=1}^{p} \sigma_{i} \rangle}{\|\nu\|^{p-\ell}} - \frac{\langle \phi, \nu^{\otimes p} \rangle}{\|\nu\|^{p}} \right\}.$$

Comparing with (7), integrating with respect to  $\zeta^{-2}\Lambda(d\zeta)$ , and putting all together, we obtain

$$\begin{aligned} \mathcal{G}_{\kappa}^{(D)}F_{\phi}(\nu) + \mathcal{G}_{\Lambda}^{(J)}F_{\phi}(\nu) &= \\ \int_{\mathcal{T}} \frac{\nu(da)}{\|\nu\|} \sum_{\ell=2}^{p} \beta_{p,\ell}^{(\Lambda)} \left( G_{\phi}^{(\ell)} \left( \frac{\nu}{\|\nu\|}; a \right) - \binom{p}{\ell} G_{\phi} \left( \frac{\nu}{\|\nu\|} \right) \right). \end{aligned}$$

This is the form of the generator of the  $\Lambda$ -Fleming-Viot process with  $\Lambda(\{0\}) = 0$ . Finally, for the diffusion term we compute

$$\begin{split} \|\nu\| F_{\phi}''(\nu; a, a) &= \|\nu\| \sum_{i=1}^{p} \frac{\|\nu\| \left(F_{\phi_{i}^{(a)}}'(\nu; a) - F_{\phi}'(\nu; a)\right) - \left(F_{\phi_{i}^{(a)}}(\nu) - F_{\phi}(\nu)\right)}{\|\nu\|^{2}} \\ &= \sum_{i=1}^{p} \left\{ F_{\phi_{i}^{(a)}}'(\nu; a) - F_{\phi}'(\nu; a) - \frac{F_{\phi_{i}^{(a)}}(\nu) - F_{\phi}(\nu)}{\|\nu\|} \right\}, \end{split}$$

where, by (33) and (34), we have

$$\int_{\mathcal{T}} \nu(da) \sum_{i=1}^{p} \left\{ F'_{\phi}(\nu; a) - \frac{F_{\phi_i^{(a)}}(\nu) - F_{\phi}(\nu)}{\|\nu\|} \right\} = 0.$$

#### REFERENCES

Thus

$$\int_{\mathcal{T}} \nu(da) \|\nu\| F_{\phi}''(\nu; a, a) = \int_{\mathcal{T}} \nu(da) \sum_{i=1}^{p} F_{\phi_{i}^{(a)}}'(\nu; a)$$

where, using (33) again, for each term in the r.h.s. above we have

$$F_{\phi_{i}^{(a)}}^{\prime}(\nu;a) = \frac{1}{\|\nu\|} \sum_{j=1}^{p} \left\{ F_{\phi_{i,j}^{(a)}}(\nu) - F_{\phi_{i}^{(a)}}(\nu) \right\}$$

where we note that  $\phi_{i,i}^{(a)} = \phi_i^{(a)}$ . Substituting above and noting also that  $\int_{\mathcal{T}} \frac{\nu(da)}{\|\nu\|} G_{\phi_i^{(a)}}(\frac{\nu}{\|\nu\|}) = G_{\phi}(\frac{\nu}{\|\nu\|}),$ 

$$\begin{split} \frac{\sigma^2}{2} \int_{\mathcal{T}} \nu(da) \left\|\nu\right\| F_{\phi}''(\nu; a, a) &= \frac{\sigma^2}{2} \int_{\mathcal{T}} \frac{\nu(da)}{\left\|\nu\right\|} \sum_{1 \le i \ne j \le p} G_{\phi_{i,j}^{(a)}}(\frac{\nu}{\left\|\nu\right\|}) - G_{\phi_{i}^{(a)}}(\frac{\nu}{\left\|\nu\right\|}) \\ &= \frac{\sigma^2}{2} \sum_{1 \le i \ne j \le p} \int_{\mathcal{T}} \frac{\nu(da)}{\left\|\nu\right\|} \left\{ G_{\phi_{i,j}^{(a)}}(\frac{\nu}{\left\|\nu\right\|}) \right\} - G_{\phi}(\frac{\nu}{\left\|\nu\right\|}). \end{split}$$

This is the form of the generator of the standard Fleming-Viot process (10). This finishes the proof of the characterization of the limiting process  $(\rho_t)_{t\geq 0}$  of the sequence  $\left(\rho_t^{(n)}\right)_{t\geq 0}$ .

Now, from the convergence of each of the coordinates in  $(\rho_t^{(n)}, \xi_t^{(n)})$  separately, we obtain  $(\rho_t^{(n)}, \xi_t^{(n)})_{t\geq 0} \xrightarrow{d} (\rho_t, \xi_t)_{t\geq 0}$  in  $D([0, \infty), \mathsf{PM}(\mathcal{T})) \times D([0, \infty), \mathbb{R})$ . Since the latter identifies the law in the limit, the corresponding convergence in  $D([0, \infty), \mathsf{PM}(\mathcal{T}) \times \mathbb{R})$  follows from the precompactness of the family of processes  $\left\{ \Phi(\left(\nu_t^{(n)}\right)_{t\geq 0})\right\}_n$  which, in turn, follows from the tightness of the family  $\left\{ \left(\nu_t^{(n)}\right)_{t\geq 0}\right\}_n$ , the continuity of the map  $\Phi(\nu) = (\nu/\|\nu\|, \log(\|\nu\|))$  on  $\mathbb{M}(\mathcal{T}) \setminus \{0\}$ , and the fact that  $(\xi_t)_{t\geq 0} \subset (-\infty, \infty)$  with probability one. Finally, the function  $\Phi^{-1}$  being continuous, another application of the mapping theorem gives the convergence of the entire sequence  $\left(\nu_t^{(n)}\right)_{t\geq 0} \xrightarrow{d} (\nu_t)_{t\geq 0}$  in  $D([0, \infty), \mathbb{M}(\mathcal{T}))$ .

By Proposition 3.3 in [18] the  $\Lambda$ -Fleming Viot process  $(\rho_t)_{t\geq 0}$  is Feller, as well as the Lévy process  $(\xi_t)_{t\geq 0}$ . The Feller property extends easily to the pair  $(\rho_t, \xi_t)_{t\geq 0}$  as in the proof of Proposition 4.4.

#### Acknowledgements

Arno Siri-Jégousse was supported by DGAPA-PAPIIT-UNAM grant IN-10282. Alejandro H. Wences was supported by the ANR LabEx CIMI (grant ANR-11-LABX-0040) within the French State Programme "Investissements d'Avenir."

### References

 Larbi Alili et al. "Inversion, duality and Doob *h*-transforms for self-similar Markov processes". In: <u>Electronic Journal of Probability</u> 22 (2017), pp. 1– 18. DOI: 10.1214/17-EJP33. URL: https://doi.org/10.1214/17-EJP33.

- Jean Bertoin. <u>Random fragmentation and coagulation processes</u>. Cambridge Studies in Advanced Mathematics. Cambridge University Press, 2006. DOI: 10.1017/CB09780511617768.
- Jean Bertoin and Igor Kortchemski. "Self-similar scaling limits of Markov chains on the positive integers". In: <u>The Annals of Applied Probability</u> 26.4 (2016), pp. 2556-2595. ISSN: 10505164. URL: http://www.jstor.org/stable/24810065.
- Jean Bertoin and Jean-François Le Gall. "Stochastic flows associated to coalescent processes". In: Probability Theory and Related Fields 126.2 (2003), pp. 261–288. ISSN: 1432-2064. DOI: 10.1007/s00440-003-0264-4. URL: https://doi.org/10.1007/s00440-003-0264-4.
- Jean Bertoin and Jean-François Le Gall. "The Bolthausen–Sznitman coalescent and the genealogy of continuous-state branching processes". In: Probability Theory and Related Fields 117 (2000), pp. 249–266. DOI: 10.1007/s004400050006.
- [6] Patrick Billingsley. <u>Convergence of probability measures</u>. 2nd ed. Wiley series in probability and statistics. Probability and statistics section. New York: Wiley, 1999. ISBN: 9780471197454.
- [7] Matthias Birkner et al. "A modified lookdown construction for the Xi-Fleming-Viot process with mutation and populations with recurrent bottlenecks". In: <u>ALEA : Latin American Journal of Probability and Mathematical Statistics</u> (2009). URL: <u>https://api.semanticscholar.org/CorpusID:2169486</u>.
- [8] Matthias Birkner et al. "Alpha-stable branching and beta-coalescents". In: <u>Electronic Journal of Probability</u> 10 (2005), pp. 303–325. DOI: 10.1214/EJP.v10-241. URL: https://doi.org/10.1214/EJP.v10-241.
- [9] Robert M. Blumenthal and Ronald K. Getoor. <u>Markov processes and potential theory</u>. Pure and Applied Mathematics. Academic Press, New York and London, 1968.
- [10] María E. Caballero, Adrián González Casanova, and José L. Pérez. "The relative frequency between two continuous-state branching processes with immigration and their genealogy". In: <u>arXiv preprint</u> (2023). DOI: https://doi.org/10.48550/arXiv.20 arXiv: 2010.00742 [math.PR].
- [11] Loïc Chaumont, Henry Pantí, and Víctor Rivero. "The Lamperti representation of real-valued self-similar Markov processes". In: <u>Bernoulli</u> 19.5B (2013), pp. 2494–2523. DOI: 10.3150/12-BEJ460. URL: https://doi.org/10.3150/12-BEJ460.
- [12] Donald A. Dawson. "Measure valued Markov processes". In: <u>Hennequin, PL. (eds) Ecole d'Eté de Probab</u> Vol. 1541. Lecture Notes in Mathematics. Springer, Berlin, Heidelberg, 1993. DOI: https://doi.org/10.1007/BFb0084190.
- [13] Donald A. Dawson and Kenneth J. Hochberg. "Wandering random measures in the Fleming-Viot model". In: <u>Ann. Probab.</u> 10.3 (1982), pp. 554–580. ISSN: 0091-1798,2168-894X. URL: http://links.jstor.org/sici?sici=0091-1798(198208)10:3%
- Peter Donnelly and Thomas G. Kurtz. "Particle representations for measure-Valued population models". In: <u>The Annals of Probability</u> 27.1 (1999), pp. 166-205. ISSN: 00911798. URL: http://www.jstor.org/stable/2652871 (visited on 02/16/2024).

- [15] Alison Etheridge and American Mathematical Society. <u>An introduction to superprocesses</u>. University lecture series. American Mathematical Society, 2000. ISBN: 9780821827062. URL: https://books.google.to/books?id=zUHyBwAAQBAJ.
- [16] Stewart N. Ethier and Thomas G. Kurtz. <u>Markov processes: characterization and convergence</u>. New York: John Wiley & Sons, 1986. ISBN: 978-0-470-31732-7.
- Patrick J. Fitzsimmons. "On the martingale problem for measure-valued Markov branching processes". In: <u>Seminar on Stochastic Processes</u>, 1991.
   Boston, MA: Birkhäuser Boston, 1992, pp. 143–156. ISBN: 978-1-4612-0381-0. DOI: 10.1007/978-1-4612-0381-0\_12. URL: https://doi.org/10.1007/978-1-4612-0381-0\_
- [18] Clément Foucart. "Generalized Fleming-Viot processes with immigration via stochastic flows of partitions". In: <u>Lat. Am. J. Probab. Math. Stat.</u> 2.9 (2012), pp. 451–472.
- [19] Alexander Gnedin, Alexander Iksanov, and Alexander Marynych. "Λcoalescents: a survey". In: Journal of Applied Probability 51A (2014), pp. 23– 40. DOI: 10.1239/jap/1417528464.
- [20] Adrián González Casanova, Imanol Nuñez, and José L. Pérez. "Alphastable branching and beta-frequency processes, beyond the IID assumption". In: <u>arXiv preprint</u> (2023). DOI: https://doi.org/10.48550/arXiv.2303.05477. arXiv: 2303.05477 [math.PR].
- [21] Adrián González Casanova et al. "Asymptotics of the frequency spectrum for general Dirichlet Xi-coalescents". In: <u>Electronic Journal of Probability</u> 29.none (2024), pp. 1–35. DOI: 10.1214/23-EJP1064. URL: https://doi.org/10.1214/23-EJP1064.
- Bénédicte Haas and Grégory Miermont. "Self-similar scaling limits of nonincreasing Markov chains". In: <u>Bernoulli</u> 17.4 (2011), pp. 1217–1247. DOI: 10.3150/10-BEJ312. URL: https://doi.org/10.3150/10-BEJ312.
- [23] Jean Jacod and Albert N. Shiryaev. Limit theorems for stochastic processes. Second. Vol. 288. Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]. Springer-Verlag, Berlin, 2003, pp. xx+661. ISBN: 3-540-43932-3. DOI: 10.1007/978-3-662-05265-5. URL: https://doi.org/10.1007/978-3-662-05265-5.
- [24] Anatole Joffe and Michel Metivier. "Weak convergence of sequences of semimartingales with applications to multitype branching processes". In: Advances in Applied Probability 18.1 (1986), pp. 20-65. ISSN: 00018678. URL: http://www.jstor.org/stable/1427238 (visited on 01/18/2024).
- Samuel G. G. Johnston and Amaury Lambert. "The coalescent structure of uniform and Poisson samples from multitype branching processes". In: <u>The Annals of Applied Probability</u> 33.6A (2023), pp. 4820–4857. DOI: 10.1214/23-AAP1934. URL: https://doi.org/10.1214/23-AAP1934.
- [26] Olav Kallenberg. <u>Foundations of modern probability</u>. eng. third. Probability Theory and Stochastic Modelling. Springer Nature Switzerland, 2021. DOI: https://doi.org/10.1007/978-3-030-61871-1.

- [27] John Kingman. "The coalescent". In: <u>Stochastic Processes and their Applications</u> 13.3 (1982), pp. 235-248. ISSN: 0304-4149. DOI: https://doi.org/10.1016/0304-4149(82)90011-4. URL: https://www.sciencedirect.com/science/article/pii/0304414982900114.
- [28] Andreas E. Kyprianou and Juan C. Pardo. "Continuous-state branching processes and self-similarity". In: Journal of Applied Probability 45.4 (2008), pp. 1140-1160. ISSN: 00219002. URL: http://www.jstor.org/stable/27596014 (visited on 11/27/2023).
- [29] Donald L. Cohn. <u>Measure theory</u>. eng. Birkhäuser Advanced Texts Basler Lehrbücher. New York, NY: Birkhäuser, 2013. ISBN: 978-1-4614-6955-1.
- [30] John Lamperti. "Semi-stable Markov processes. I". In: Zeitschrift für Wahrscheinlichkeitstheorie und Verv 22.3 (1972), pp. 205–225. ISSN: 1432-2064. DOI: 10.1007/BF00536091. URL: https://doi.org/10.1007/BF00536091.
- [31] Zenghu Li. Measure-valued branching markov processes. Series Title Probability Theory and Stochastic Modelling. Springer Berlin, Heidelberg, 2022. ISBN: 978-3-662-66910-5. DOI: https://doi.org/10.1007/978-3-662-66910-5.
- [32] Edwin A. Perkins. "Conditional Dawson-Watanabe processes and Fleming-Viot processes". In: <u>Seminar on Stochastic Processes</u>, 1991. Boston, MA: Birkhäuser Boston, 1992, pp. 143–156. ISBN: 978-1-4612-0381-0. DOI: 10.1007/978-1-4612-0381-0\_12. URL: https://doi.org/10.1007/978-1-4612-0381-0\_12.
- [33] Edwin A. Perkins, Erwin Bolthausen, and Aad Vaart. Part II: Dawson-Watanabe superprocesses and mea Apr. 2004, pp. 125–329.
- [34] Jim Pitman. "Coalescents with multiple collisions". In: <u>The Annals of Probability</u> 27.4 (1999), pp. 1870–1902. DOI: 10.1214/aop/1022874819. URL: https://doi.org/10.1214/aop/10228
- [35] Serik Sagitov. "The general coalescent with asynchronous mergers of ancestral lines". In: Journal of Applied Probability 36.4 (1999), pp. 1116– 1125. DOI: 10.1239/jap/1032374759.
- [36] Stanley A. Sawyer. "A formula for semigroups, with an application to branching diffusion processes". In: <u>Transactions of the American Mathematical Society</u> 152.1 (1970), pp. 1–38. ISSN: 00029947. URL: http://www.jstor.org/stable/1995636 (visited on 10/15/2023).
- [37] Jason Schweinsberg. "Coalescent processes obtained from supercritical Galton-Watson processes". In: <u>Stoch. Proc. Their Appl.</u> 106.1 (2003), pp. 107– 139. ISSN: 0304-4149. DOI: https://doi.org/10.1016/S0304-4149(03)00028-0. URL: https://www.sciencedirect.com/science/article/pii/S0304414903000280.